Note: See sections 5 and 6.

Permissible	e ingredients and requirements		
Column 1	Column 2	Column 3	Column 4
Item	Ingredient Name	Purpose	Specific requirements
2855	KADSURA COCCINEA	A, H	
2856	KAEMPFERIA GALANGA	А, Н	
2857	KALMIA LATIFOLIA	А, Н	Beta-arbutin is a mandatory component of Kalmia latifolia.
			When for oral use, the maximum recommended daily dose must not provide more than 500 mg of beta- arbutin.
			When for dermal application exclusively to the face:
			a) the concentration of beta-arbutin in the medicine must not be more than 7%;
			b) hydroquinone is a mandatory component; and
			c) the concentration of hydroquinone must not be more than 10 mg/kg or 10 mg/L or 0.001%.
			When for use other than oral or dermal application exclusively to the face, the concentration of beta-arbutin in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%.
2858	KAOLIN	Е	
2859	KELP DRY	А, Н	Iodine is a mandatory component of Kelp dry.
			Only for external use when the concentration of iodine in the medicine (excluding salts derivatives or iodophors) is 2.5% or less.
			Only for internal use when the medicine contains less than 300 micrograms of

Volume 4	4
----------	---

			iodine per maximum recommended daily dose.
2860	KELP POWDER	A, E, H	Iodine is a mandatory component of Kelp powder.Only for external use when the concentration of iodine in the medicine (excluding salts derivatives or iodophors) is 2.5% or less.Only for internal use when the medicine contains less than 300 micrograms of iodine per maximum recommended daily dose.
2861	KERATIN	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 5%.
2862	KEROSENE	E, H	Only for use as a homoeopathic ingredient. When used in liquid preparations, the concentration in the medicine must be no more than 25%.
2863	KHAYA SENEGALENSIS	A, E	<ul> <li>The maximum daily dose of the medicine must not contain more than the equivalent of 1 g dry bark of Khaya senegalensis.</li> <li>The following warning statements are required on the medicine label: <ul> <li>(PREGNT2) 'Do not use if pregnant or likely to become pregnant' (or words to that effect);</li> <li>(LONGUSE) 'Not for prolonged use. May harm liver';</li> <li>(GEN2) 'If symptoms persist, seek the advice of a healthcare professional';</li> <li>(CHILD3) 'Use in children under 12 years is not recommended'; and</li> <li>(7DAYS) 'Do not use for more than 7 days'.</li> </ul> </li> </ul>

2864	KIDNEY BEAN	Е	
2865	KIRSCH	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
2866	KIWI FRUIT	E	
2867	KNAUTIA ARVENSIS	A, H	
2868	KOREAN GINSENG ROOT DRY	A, H	
2869	KOREAN GINSENG ROOT POWDER	A, H	
2870	KRAMERIA IXINE	A, H	
2871	KRAMERIA LAPPACEA	A, H	
2872	KUNZEA AMBIGUA	A	Only for use when the plant preparation is essential oil.
			Only for use when the route of administration is topical or inhalation.
			When the dosage form is essential oil, a restricted flow insert must be fitted on the container and the medicine requires the following warning statements on the medicine label:
			- (CHILD) 'Keep out of reach of children'
			- (EXTERN) 'For external use only'
			- (UNDILU) 'Not to be applied undiluted to the skin except on the advice of a health care practitioner'.
			When the dosage form is other than essential oil, the maximum concentration in topical medicines must be no more than 25% w/w and the medicine requires the following warning statements on the medicine label:
			- (CHILD) 'Keep out of reach of children'
			- (EXTERN) 'For external use only'.

Volume	4
--------	---

2873	L-BORNEOL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
2874	L-BORNYL ACETATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used as in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
2875	L-CARVONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
2876	L-LIMONENE	E	L-limonene must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation or a fragrance proprietary excipient formulation. The total concentration of flavour proprietary excipient formulations
			<ul> <li>proprietary excipient formulations</li> <li>containing l-limonene must not be more than 5% of the total medicine.</li> <li>The total concentration of fragrance</li> <li>proprietary excipient formulations</li> <li>containing l-limonene must not be more than 1% of the total medicine.</li> </ul>

2877	L-LINALOOL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
2878	L-MENTHONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
2879	L-MENTHYL ACETATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
2880	L-ROSE OXIDE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
2881	LABDANUM ABSOLUTE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li> </ul>

### Volume 4

2882	LABDANUM GUM EXTRACT ETHYL ESTER	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance and the total fragrance concentration in a medicine is no more than 1%.</li></ul>
2883	LABDANUM OIL	A, E, H	
2884	LABURNUM ANAGYROIDES	A, H	Sparteine is a mandatory component of Laburnum anagyroides. The concentration of sparteine in the medicine must be no more than 0.001%.
2885	LACTALBUMIN	E	
2886	LACTIC ACID	А, Е, Н	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing form time to time.Sponsors should consider the impact of excipients containing alpha hydroxy acids on the sensitivity of the skin to sunlight and should ensure the finished medicine is safe for its intended purpose
2887	LACTITOL	E	
2888	LACTITOL MONOHYDRATE	Е	
2889	LACTO-N-NEOTETRAOSE	A	<ul> <li>Lactose is a mandatory component of lacto-N-neotetraose.</li> <li>The route of administration for medicines that contain lacto-N-neotetraose must be limited to oral.</li> <li>The maximum recommended daily dose of the medicine must not provide more than:</li> <li>(a) 1.5 g of lacto-N-neotetraose to individuals aged 4 years and older; and</li> <li>(b) 0.6 g of lacto-N-neotetraose to individuals aged up to 3 years (inclusive).</li> </ul>

			<ul> <li>One of the following statements (or words to the same effect) is required on the medicine label:</li> <li>(a) When the medicine is only for use in individuals aged 2 years and above: 'Not to be taken on the same day with other products containing lacto-N-neotetraose'; or</li> <li>(b) When the medicine is for use in children aged less than 2 years: 'Not to be taken on the same day with breastmilk or other products containing lacto-N-neotetraose'.</li> </ul>
2890	LACTO-N-TETRAOSE	A	Lactose is a mandatory component of lacto-N-tetraose. The route of administration for medicines that contain lacto-N-tetraose
			must be limited to oral. The maximum recommended daily dose of the medicine must not provide more than:
			a) 2 g of lacto-N-tetraose to individuals aged 1 year and older; and
			b) 0.6 g of lacto-N-tetraose to individuals aged more than 6 months to 11 months (inclusive); and
			c) 0.8 g of lacto-N-tetraose to individuals aged up to 6 months (inclusive).
			One of the following statements (or words to the same effect) is required on the medicine label:
			a) When the medicine is only for use in individuals aged 2 years and above: 'Not to be taken on the same day with other products containing lacto-N-tetraose'; or
			b) When the medicine is for use in children aged less than 2 years: 'Not to be taken on the same day with breastmilk or other products containing lacto-N- tetraose'.
2891	LACTOBACILLUS ACIDOPHILUS	A	

Vol	lume	4

2892	LACTOBACILLUS AMYLOVORUS	А	
2893	LACTOBACILLUS BREVIS	А	
2894	LACTOBACILLUS CASEI	А	
2895	LACTOBACILLUS CASEI SUBSP. BIOVAR CASEI	А	
2896	LACTOBACILLUS CRISPATUS	Α	
2897	LACTOBACILLUS DELBRUECKII SSP BULGARICUS	А	
2898	LACTOBACILLUS DELBRUECKII SSP LACTIS	А	
2899	LACTOBACILLUS FERMENTUM	А	
2900	LACTOBACILLUS GALLINARUM	А	
2901	LACTOBACILLUS GASSERI	А	
2902	LACTOBACILLUS HELVETICUS	A	
2903	LACTOBACILLUS JOHNSONII	А	
2904	LACTOBACILLUS KEFIRANOFACIENS	A	
2905	LACTOBACILLUS KEFIRGRANUM	А	
2906	LACTOBACILLUS KEFIRI	А	
2907	LACTOBACILLUS PARACASEI	А	
2908	LACTOBACILLUS PARACASEI SUBSP. PARACASEI	A	
2909	LACTOBACILLUS PLANTARUM	А	
2910	LACTOBACILLUS REUTERI	А	
2911	LACTOBACILLUS RHAMNOSUS	А	
2912	LACTOBACILLUS SALIVARIUS SSP SALICINIUS	A	
2913	LACTOBACILLUS SALIVARIUS SSP SALIVARIUS	А	
2914	LACTOBIONIC ACID	Е	Only for use in topical medicines for dermal application.
2915	LACTOSCATONE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
2916	LACTOSE	Е	

## Volume 4

2917	LACTOSE MONOHYDRATE	Е	
2918	LACTUCA SATIVA	А, Н	
2919	LACTUCA VIROSA	A, H	
2920	LACTULOSE	Е	
2921	LACTULOSE SOLUTION	A	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force on existing form time to time.
2922	LAGENARIA VULGARIS	A, H	
2923	LAMINARIA CLOUSTONI	A, E, H	Iodine is a mandatory component of Laminaria cloustoni.
			Only for external use when the concentration of iodine in the medicine (excluding salts derivatives or iodophors is 2.5% or less.
			Only for internal use when the medicine contains less than 300 micrograms of iodine per maximum recommended daily dose.
2924	LAMINARIA DIGITATA	A, E, H	Iodine is a mandatory component of Laminaria digitata.
			Only for external use when the concentration of iodine in the medicine (excluding salts derivatives or iodophors is 2.5% or less.
			Only for internal use when the medicine contains less than 300 micrograms of iodine per maximum recommended daily dose.
2925	LAMINARIA JAPONICA	A, E, H	Iodine is a mandatory component of Laminaria japonica.
			Only for external use when the concentration of iodine in the
			medicine (excluding salts derivatives or iodophors) is 2.5% or less.
			Only for internal use when the medicine contains less than 300 micrograms of

### Volume 4

			iodine per maximum recommended daily dose.
2926	LAMIUM ALBUM	A, H	
2927	LANETH-5	E	Only for use in topical medicines for dermal application.
2928	LANOLIN ALCOHOL	E	Only for use in topical medicines for dermal application.
2929	LANOLIN OIL	E	Only for use in topical medicines for dermal application.
2930	LANOLIN WAX	E	Only for use in topical medicines for dermal application.
2931	LANTANA CAMARA	A, H	The maximum recommended daily dose must contain no more than 1mg of the equivalent dry herbal material of Lantana camara.
2932	LARIX ARABINOGALACTAN	A, E	<ul> <li>The concentration of polysaccharides in the ingredient must be greater than or equal to 85%.</li> <li>The ingredient must be derived from Larix occidentalis or Larix larcinia.</li> <li>Only for use in oral medicines or topical medicines for dermal application, and not to be included in topical products intended for use in the eye.</li> <li>The maximum recommended daily dose of Larix arabinogalactan in oral medicines must not be more than 15 grams.</li> <li>The concentration of Larix arabinogalactan in topical medicines for dermal applications for dermal application must not exceed 5.0%.</li> </ul>
2933	LARIX DECIDUA	A, H	
2934	LARIX KAEMPFERI	А, Н	The maximum recommended daily dose must be no more than 1 mg of the

			equivalent dry herbal material of Larix kaempferi.
2935	LARREA TRIDENTATA	A, H	The following warning statement is required on the medicine label:
			(CHAP1) 'In rare cases, Larrea tridentata may harm the liver. Stop use and see a doctor if you have yellowing skin/eyes o unusual: fatigue, nausea, appetite loss, abdominal pain or dark urine.'
2936	LATHYRUS SATIVUS	А, Н	The maximum recommended daily dose must be no more than 1mg of the equivalent dry herbal material of Lathyrus sativus.
			The medicine must not contain lathyrogenic amino acids.
2937	LAURAMINE OXIDE	E	
2938	LAUREL LEAF OIL	A, H	When the total concentration of bay oil in the medicine is more than 25%:
			(a) the nominal capacity of the container must not be more than 25 mL;
			(b) the container must be fitted with a restricted flow insert;
			(c) the following warning statements are required on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect);
			- (NTAKEN) 'Not to be taken'; and
			(d) when the nominal capacity of the container is greater than 15 mL, the container must also be fitted with a child resistant closure.
2939	LAURETH-10	E	Only for use in topical medicines for dermal application.
2940	LAURETH-12	E	Only for use in topical medicines for dermal application.

Volume -	4
----------	---

2941	LAURETH-2	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.4%.
			Residual levels of ethylene oxide (and related substances) must be kept below the level of detection.
2942	LAURETH-23	E	Only for use in topical medicines for dermal application.
2943	LAURETH-3	Е	Only for use in topical medicines for dermal application.
2944	LAURETH-4	Е	Only for use in topical medicines for dermal application.
2945	LAURETH-7	Е	Only for use in topical medicines for dermal application.
2946	LAURETH-8	Е	
2947	LAURIC ACID	A, E	When for use as an active ingredient is for use in oral medicines only and the maximum recommended daily dose must not exceed 1500 mg.
2948	LAURIL MACROGOL 400 DIMETICONE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or on damaged skin.
			The concentration in the medicine must be no more than 5%.
2949	LAUROMACROGOL 400	E	Only for use in topical medicines for dermal application.
2950	LAUROYL LYSINE	E	Only for use in topical medicines for dermal application and not to be included in medicines for use in the eye or on damaged skin. The concentration in the medicine must be no more than 5.0%.

2951	LAURUS NOBILIS	A, E, H	When the plant preparation is oil or distillate, and the concentration of bay oil or distillate in the medicine is greater than 25%:
			(a) the nominal capacity of the container must not be more than 25 mL;
			(b) the container must be fitted with a restricted flow insert;
			(c) the following warning statements are required on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect);
			- (NTAKEN) 'Not to be taken'; and
			(d) when the nominal capacity of the container is greater than 15 mL, the container must also be fitted with a child resistant closure.
2952	LAURYL ALDEHYDE	E	Permitted for use only in combination with other permitted ingredients as a coating solution, flavour or a fragrance. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
2953	LAURYL BETAINE	Е	Only for use in topical medicines for dermal application.
2954	LAURYL GLUCOSIDE	E	Only for use as an excipient ingredient in topical medicines for dermal application and not to be included in topical medicines intended for use in the eye. The concentration in the medicine must be no more than 12%.
2955	LAURYL LACTATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.

Volume 4

			The concentration in the medicine must be no more than 3%. Sponsors should consider the impact of excipients containing alpha hydroxy acids on the sensitivity of the skin to sunlight and should ensure the finished medicine is safe for its intended purpose.
2956	LAURYL PCA	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 1%.
2957	LAURYL PEG-10 TRIS(TRIMETHYLSILOXY)SILYLETH YL DIMETICONE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or on damaged skin. The concentration in the medicine must be no more than 2%.
2958	LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETICONE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or damaged skin. The concentration in the medicine must be no more than 3.5%.
2959	LAURYL PEG/PPG-18/18 METHICONE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 9%. Residual levels of ethylene oxide (and related substances) must be kept below the level of detection.
2960	LAURYL POLYGLUCOSE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration must not exceed 1% in leave-on medicines and 3% in wash- on/wash-off medicines.

15

2961	LAURYL PYRROLIDONE	E	Only for use in topical medicines for dermal application.
2962	LAURYLDIMONIUM HYDROXYPROPYL HYDROLYSED COLLAGEN	E	Only for use in topical medicines for dermal application.
2963	LAURYLDIMONIUM HYDROXYPROPYL HYDROLYSED SOY PROTEIN	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.007%.
2964	LAURYLMETICONE COPOLYOL	E	Only for use in topical medicines for dermal application.
2965	LAVANDIN OIL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
2966	LAVANDIN OIL ABRIAL	A, E, H	
2967	LAVANDIN OIL GROSSO	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
2968	LAVANDULA ANGUSTIFOLIA	A, E, H	<ul> <li>Camphor is a mandatory component of Lavandula angustifolia.</li> <li>In solid and semi solid preparations, the concentration of camphor must be no more than 12.5%.</li> <li>In liquid preparations other than essential oils or distillates, the concentration of camphor must be no more than 2.5%.</li> </ul>

Volume	4

2969	LAVANDULA ANGUSTIFOLIA SUBSP. ANGUSTIFOLIA	A, E, H	Camphor is a mandatory component of Lavandula angustifolia subsp. angustifolia.
			In solid and semi solid preparations, the concentration of camphor must be no more than 12.5%.
			In liquid preparations other than essential oils or distillates, the concentration of camphor must be no more than 2.5%.
2970	LAVANDULA X INTERMEDIA	A, E, H	Camphor is a mandatory component of Lavandula x intermedia.
			In solid and semi solid preparations, the concentration of camphor must be no more than 12.5%.
2971	LAVENDER OIL	A, E, H	
2972	LAWSONIA INERMIS	A, H	
2973	LEAD	Н	Only for use as an active homoeopathic ingredient.
			The concentration in the medicine must be no more than 0.001%.
2974	LEAD ACETATE	Н	Only for use as an active homoeopathic ingredient.
2975	LEAF ACETAL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used as a flavour the total flavour concentration in a medicine must be no more than 5%.
2976	LECITHIN	A, E	
2977	LEDEBOURIELLA SESELOIDES	A, H	
2978	LEDUM PALUSTRE	A, H	Beta-arbutin is a mandatory component of Ledum palustre.
			When for dermal application exclusively to the face:
			a) the concentration of beta-arbutin in the medicine must not be more than 7%;

			b) hydroquinone is a mandatory component; and
			c) the concentration of hydroquinone must not be more than 10 mg/kg or 10 mg/L or 0.001%.
			When for topical use other than dermal application exclusively to the face, the concentration of beta-arbutin in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%. When the route of administration is other than topical, the maximum recommended daily dose must not contain more than 0.001 mg of the equivalent dry herbal material of Ledum palustre.
2979	LEMNA MINOR	A, H	
2980	LEMON	E	When used internally, oxedrine is a mandatory component of lemon. The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
2981	LEMON BALM LEAF DRY	A, H	
2982	LEMON BALM LEAF POWDER	A, E, H	
2983	LEMON OIL	A, E, H	When used internally, oxedrine is a mandatory component of lemon oil. The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams. The warning statement (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect) must be included on the medicine label unless the medicine is: a) steam distilled or rectified; or b) for internal use; or c) contains 0.05% or less of lemon oil; or d) for use in soaps or bath or shower gels that are washed off the skin.

Volume 4

Volume	4
--------	---

2984	LEMON OIL DISTILLED	A, E, H	When used internally, oxedrine is a mandatory component of lemon oil distilled.
			The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
2985	LEMON OIL TERPENELESS	A, E, H	When used internally, oxedrine is a mandatory component of lemon oil terpeneless.
			The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
2986	LEMON OIL TERPENES AND TERPENOIDS	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
2987	LEMON PEEL DRIED	A, E, H	When used internally, oxedrine is a mandatory component of lemon peel dried.
			The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
2988	LEMONGRASS OIL	A, E, H	
2989	LENS CULINARIS	A, H	
2990	LENTIL	E	
2991	LENTINULA EDODES	A, E, H	
2992	LEONTOPODIUM ALPINUM	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must
2993	LEONURUS CARDIACA	A, E, H	be no more than 1%.

Volume 4	Vol	lume	4
----------	-----	------	---

2994	LEONURUS SIBIRICUS	А, Е, Н	
2995	LEPIDIUM APETALUM	A, H	
2996	LEPIDIUM MEYENII	A	<ul> <li>The route of administration for medicines that contain Lepidium meyeni must be limited to oral.</li> <li>The ingredient must consist of the dried tuber of Lepidium meyenii only.</li> <li>The maximum recommended daily dose of the medicine must not provide more than 3.5 g of Lepidium meyenii dried tuber (or its extract equivalent).</li> </ul>
2997	LEPTOSPERMUM PETERSONII	E	Only for use in topical medicines for dermal application. The concentration in the medicine must be no more 5%.
2998	LEPTOSPERMUM SCOPARIUM OIL	A	<ul> <li>Only for use as an active ingredient when the route of administration is topical or oral application in a mouthwash preparation.</li> <li>If the concentration is more than 25%, the nominal capacity of the container must be no more than 25mL.</li> <li>When the concentration is more than 25%, and the nominal capacity of the container less than 15mL, a restricted flow insert must be fitted on the container and requires the following warning statements on the medicine label: <ul> <li>(CHILD) 'Keep out of reach of children' (or word to that effect)</li> <li>(NTAKEN) 'Not to be taken'</li> </ul> </li> <li>When the concentration is more than 25%, the nominal capacity of the container is more than 15 mL but no more than 25 mL, a child resistant closure and restricted flow insert must be fitted on the container and requires the following warning statements on the medicine label:</li> </ul>

#### Volume 4

			<ul><li> (CHILD) 'Keep out of reach of children' (or word to that effect)</li><li> (NTAKEN) 'Not to be taken'</li></ul>
2999	LESPEDEZA CAPITATA	A, H	
3000	LETTUCE	Е	
3001	LEUCINE	A, E	
3002	LEUZEA UNIFLORA	A, H	
3003	LEVISTICUM OFFICINALE	A, H	
3004	LEVOCARNITINE	А	
3005	LEVOCARNITINE FUMARATE	А	
3006	LEVOCARNITINE HYDROCHLORIDE	А	
3007	LEVOCARNITINE MAGNESIUM CITRATE	А	
3008	LEVOCARNITINE TARTRATE	A	
3009	LEVOMEFOLATE CALCIUM	A	<ul> <li>Available for medicines intended for internal use only.</li> <li>Levomefolic acid is a mandatory component of levomefolate calcium.</li> <li>The maximum recommended daily dose must not provide more than 500 micrograms of levomefolic acid from levomefolate calcium.</li> <li>When the medicine contains a combination of folic acid, folinic acid or levomefolic acid, the medicine must not provide more than a combined total of 500 micrograms of folic acid, folinic acid and levomefolic acid per maximum recommended daily dose.</li> </ul>
3010	LEVOMEFOLATE GLUCOSAMINE	A	<ul> <li>Available for medicines intended for internal use only.</li> <li>Levomefolic acid is a mandatory component of levomefolate glucosamine.</li> <li>The maximum recommended daily dose must not provide more than 500 micrograms of levomefolic acid from levomefolate glucosamine.</li> <li>When the medicine contains a combination of folic acid, folinic acid or levomefolic acid, the medicine must not</li> </ul>

			Volume 4
			provide more than a combined total of 500 micrograms of folic acid, folinic acid and levomefolic acid per maximum recommended daily dose.
3011	LEVOTHYROXINE SODIUM	Н	Only for use as an active homoeopathic ingredient.
3012	LEVULINIC ACID	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3013	LIGHT KAOLIN	E	
3014	LIGHT LIQUID PARAFFIN	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.
3015	LIGHT MAGNESIUM OXIDE	A, E, H	Magnesium is a mandatory component of light magnesium oxide.
			When used in a medicine:
			<ul><li>(a) with an oral route of administration;</li><li>(b) not indicated for laxative (or related) use; and</li></ul>
			<ul><li>(c) where the maximum recommended daily dose for:</li></ul>
			<ul> <li>(i) children aged between 1 and 3 years</li> <li>(inclusive) provides 65 mg or more total</li> <li>magnesium from inorganic magnesium</li> <li>salts;</li> </ul>
			<ul><li>(ii) children aged between 4 and 8 years</li><li>(inclusive) provides 110 mg or more</li><li>total magnesium from inorganic</li><li>magnesium salts; or</li></ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total

Volume	4

		magnesium from inorganic magnesium salts;
		the following warning statement is required on the medicine label:
		- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
		When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.
LIGUSTICUM SINENSE	A, H	
LIGUSTICUM STRIATUM	A, E, H	
LIGUSTRUM LUCIDUM	A, H	
LILIUM BROWNII	A, H	
LILIUM CANDIDUM	A, E, H	
LILIUM LANCIFOLIUM	A, H	
LILIUM LONGIFLORUM	A, H	
LIME FRUIT	Е	
LIME OIL	E	<ul> <li>Lime oil must only be included in medicines when in combination with other permitted ingredients as a flavour or a fragrance proprietary excipient formulation.</li> <li>The total concentration of flavour proprietary excipient formulations containing lime oil must not be more than 5% of the total medicine.</li> <li>The total concentration of fragrance proprietary excipient formulations containing lime oil must not be more than 5% of the total medicine.</li> <li>The total concentration of fragrance proprietary excipient formulations containing lime oil must not be more than 1% of the total medicine.</li> <li>When for other than internal use: <ul> <li>(a) the concentration of lime oil in the medicine must not be more than 0.5%; o</li> <li>(b) the following warning statement (or words to the same effect) is required on the medicine label:</li> </ul> </li> </ul>
	LIGUSTICUM STRIATUMLIGUSTRUM LUCIDUMLILIUM BROWNIILILIUM CANDIDUMLILIUM LANCIFOLIUMLILIUM LONGIFLORUMLIME FRUIT	LIGUSTICUM STRIATUMA, E, HLIGUSTRUM LUCIDUMA, HLILIUM BROWNIIA, HLILIUM CANDIDUMA, E, HLILIUM LANCIFOLIUMA, HLILIUM LONGIFLORUMA, HLIME FRUITE

			v olulile -
3025	LIME OIL COLDPRESSED	A, E, H	The warning statement (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect) must be included on the medicine label unless the medicine is: a) for internal use; or b) contains 0.5% or less of lime oil coldpressed; or c) for use in soaps or bath or shower gels that are washed off the skin.
3026	LIME OIL DISTILLED	A, E, H	<ul> <li>The warning statement (SENS)</li> <li>'Application to skin may increase sensitivity to sunlight' (or words to that effect) must be included on the medicine label unless the medicine is:</li> <li>a) for internal use; or</li> <li>b) contains 0.5% or less of lime oil distilled; or</li> <li>c) for use in soaps or bath or shower gels that are washed off the skin.</li> </ul>
3027	LIME OIL TERPENELESS	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3028	LIME OIL TERPENES AND TERPENOIDS	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3029	LIME TREE FLOWER DRY	A, H	
3030	LIME TREE FLOWER POWDER	A, H	
3031	LIME, ESSENCE	E	

Volume 4

3032	LIMES TERPENES	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3033	LIMONENE	E	When for oral use, the quantity must be no more than 10 mg per maximum recommended daily dose.
3034	LINALOOL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3035	LINALOOL OXIDE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3036	LINALYL ACETAL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3037	LINALYL ACETATE	E	Permitted for use only:

			v olulie 4
			(a) in topical medicines for dermal application; and
			(b) in oral medicines in combination with other permitted ingredients as part of a flavour proprietary excipient formulation.
			When used in a flavour, the total flavour proprietary excipient formulation in a medicine must be no more than 5%.
3038	LINALYL BENZOATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3039	LINALYL BUTYRATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3040	LINALYL CINNAMATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3041	LINALYL FORMATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>

Volume	4
--------	---

3042	LINALYL ISOBUTYRATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3043	LINALYL PROPIONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3044	LINDERA STRYCHNIFOLIA	A, H	
3045	LINOLEAMIDOPROPYL PG- DIMONIUM CHLORIDE PHOSPHATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.5%.
3046	LINOLEIC ACID	E	
3047	LINOLENIC ACID	Е	
3048	LINSEED DRY	A, E, H	
3049	LINSEED OIL	A, E, H	
3050	LINSEED OIL FATTY ACIDS	E	Linseed oil fatty acids must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation. The total concentration of flavour proprietary excipient formulations containing linseed oil fatty acids must not be more than 5% of the total medicine.

Volume	4
--------	---

3051	LINSEED POWDER	А, Е, Н	
3052	LINUM USITATISSIMUM	A, E, H	
3053	LIPASE	A	Permitted for use only when derived from Rhizopus oryzae and in medicines containing 20,000 lipase units (equivalent to 20,000 BP units) or less of lipase activity per dosage unit. Lipase must comply with the relevant compositional guideline.
3054	LIPPIA DULCIS	A, H	
3055	LIQUID GLUCOSE	Е	
3056	LIQUID PARAFFIN	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.
3057	LIQUIDAMBAR FORMOSANA	A, H	
3058	LIQUIDAMBAR ORIENTALIS	A, H	
3059	LIQUIDAMBAR STYRACIFLUA	A, E, H	
3060	LIQUIDAMBAR STYRACIFLUA RESIN	Ε	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3061	LIQUIDAMBAR TAIWANIANA	A, H	
3062	LIQUORICE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3063	LIQUORICE DRY	A, E, H	
3064	LIQUORICE LIQUID EXTRACT	A, E, H	
3065	LIQUORICE POWDER	A, E, H	

Volume	4
--------	---

3066	LITCHI CHINENSIS	A, E, H	When used as an excipient, Litchi chinensis must only be included in medicines when the plant part is fruit, in combination with other permitted ingredients as a flavour proprietary excipient formulation.
			The total concentration of flavour proprietary excipient formulations containing Litchi chinensis must not be more than 5% of the total medicine.
3067	LITHIUM CARBONATE	Н	Only for use as an active homoeopathic ingredient.
3068	LITHOSPERMUM OFFICINALE	A, H	The maximum recommended daily dose must be no more than 1mg of the equivalent dry herbal material of Lithospermum officinale.
3069	LITSEA CUBEBA	A, E, H	
3070	LITSEA CUBEBA OIL	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3071	LOBARIA PULMONARIA	A, H	
3072	LOBELIA DRY	A, H	The concentration in the medicine must be no more than 0.001% or 10mg/kg or 10ml/L or 10 ppm unless the medicine is administered by inhalation.
3073	LOBELIA INFLATA	A, H	The concentration in the medicine must be no more than 0.001% or 10mg/kg or 10ml/L or 10 ppm unless the medicine is administered by inhalation.
3074	LOBELIA POWDER	A, H	The concentration in the medicine must be no more than 0.001% or 10mg/kg or

			10ml/L or 10 ppm unless the medicine is administered by inhalation.
3075	LOLIUM PERENNE	A, H	
3076	LOLIUM TEMULENTUM	A, H	
3077	LONGIFOLENE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total longifolene concentration in a medicine must be no more than 1%.
3078	LONICERA CAPRIFOLIUM	A, E, H	
3079	LONICERA JAPONICA	A, E, H	
3080	LONICERA PERICLYMENUM	A, H	
3081	LOPHATHERUM GRACILE	A, H	
3082	LOQUAT	Е	
3083	LORANTHUS PARASITICUS	A, H	
3084	LOROPETALUM CHINENSE	A, H	
3085	LOTUS CORNICULATUS	A, H	
3086	LOVAGE OIL	A, E, H	
3087	LOVAGE ROOT DRY	A, H	
3088	LOVAGE ROOT POWDER	A, H	
3089	LUDWIGIA PROSTRATA	A, H	
3090	LUFFA CYLINDRICA	A, H	
3091	LUFFA PURGANS	A, H	
3092	LUTEIN	A, E, H	When used as an excipient, permitted for use only as a colour in medicines limited to topical and oral routes of administration.
3093	LYCHEE	Е	
3094	LYCIUM BARBARUM	A, H	
3095	LYCIUM CHINENSE	A, E, H	
3096	LYCOPENE	A, E	
3097	LYCOPERSICON ESCULENTUM	A, E, H	Steroidal alkaloids calculated as solanine is a mandatory component of Lycopersicon esculentum.

### Volume 4

			The maximum daily dose must not provide more than 10 mg of steroidal alkaloids calculated as solanine.
3098	LYCOPODIUM ANNOTINUM	A, H	
3099	LYCOPODIUM CLAVATUM	A, H	
3100	LYCOPODIUM COMPLANATUM	A, H	
3101	LYCOPUS EUROPAEUS	A, H	
3102	LYCOPUS LUCIDUS	A, H	
3103	LYCOPUS VIRGINICUS	А, Н	Pulegone is a mandatory component of Lycopus virginicus. The concentration of pulegone in the medicine must be no more than 4%.
3104	LYGODIUM JAPONICUM	A, H	
3105	LYSIMACHIA CHRISTINAE	A, H	
3106	LYSIMACHIA VULGARIS	A, H	
3107	LYSINE	A, E	
3108	LYSINE HYDROCHLORIDE	A, E	
3109	LYTHRUM HYSSOPIFOLIA	A, H	
3110	LYTHRUM SALICARIA	A, H	
3111	LYTHRUM VERTICILLATUM	A, H	
3112	MACADAMIA INTEGRIFOLIA	A, E	
3113	MACADAMIA NUT OIL	Е	
3114	MACADAMIA TERNIFOLIA	A, E, H	
3115	MACE	E	<ul> <li>Safrole is a mandatory component of Mace.</li> <li>When used internally, the concentration of safrole in the medicine must be no more than 0.1%.</li> <li>When used topically, the concentration of safrole in the medicine must be no more than 1.0%.</li> </ul>
3116	MACE OIL	A, H	Safrole is a mandatory component of Mace oil. When used internally, the concentration of safrole in the medicine must be no more than 0.1%.

			<ul> <li>When used topically, the concentration of safrole in the medicine must be no more than 1.0%.</li> <li>When the concentration of mace oil in the preparation is more than 50% and the nominal capacity of the container is 25 mL or less, a restricted flow insert must be fitted on the container.</li> </ul>
3117	MACROCYSTIS PYRIFERA	A, E, H	Iodine is a mandatory component of Macrocystis pyrifera.Only for external use when the concentration of iodine in the medicine (excluding salts derivatives or iodophors is 2.5% or less.Only for internal use when the medicine
			contains less than 300 micrograms of iodine per maximum recommended daily dose.
3118	MACROGOL 1000	E	
3119	MACROGOL 1450	Е	Only for use in topical medicines for dermal application.
3120	MACROGOL 1500	E	
3121	MACROGOL 1500 CASTOR OIL	E	Only for use in topical medicines for dermal application and not to be include in medicines intended for use in the eye or on damaged skin. The concentration in the medicine must
			be no more than 2%.
3122	MACROGOL 200	E	Only for use in topical medicines for dermal application.
3123	MACROGOL 20000	E	
3124	MACROGOL 300	E	
3125	MACROGOL 3000	E	
3126	MACROGOL 3350	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an

Volume 4

			uncompounded substance monograph of the British Pharmacopoeia, as in force or existing form time to time.
3127	MACROGOL 40	E	Only for use in topical medicines for dermal application.
3128	MACROGOL 400	E	
3129	MACROGOL 4000	Е	
3130	MACROGOL 45000	E	Only for use in topical medicines for dermal application.
3131	MACROGOL 600	E	
3132	MACROGOL 6000	Е	
3133	MACROGOL 600000	Е	
3134	MACROGOL 800	Е	
3135	MACROGOL 8000	Е	
3136	MACROGOL 900	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.95%.
3137	MACROGOL POLY(VINYL ALCOHOL) GRAFTED POLYMER	E	Only for use in oral medicines. The concentration in the medicine must be no more than 5%.
3138	MACROPIPER EXCELSUM VAR EXCELSUM	A, H	
3139	MAGNESIUM AMINO ACID CHELATE	A, E, H	Only for use in oral medicines. The concentration of magnesium must be no more than 25% of the magnesium amino acid chelate.
3140	MAGNESIUM ASCORBATE	A, E, H	
3141	MAGNESIUM ASCORBATE MONOHYDRATE	A, E, H	
3142	MAGNESIUM ASCORBYL PHOSPHATE	Е	Only for use in topical medicines for dermal application.
3143	MAGNESIUM ASPARTATE	A, E, H	

3144	MAGNESIUM ASPARTATE DIHYDRATE	A, E, H	
3145	MAGNESIUM ASPARTATE TETRAHYDRATE	A, E, H	
3146	MAGNESIUM CARBONATE HYDRATE	A, E, H	
3147	MAGNESIUM CHLORIDE 4.5- HYDRATE	A	Magnesium is a mandatory component of magnesium chloride 4.5-hydrate. When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			<ul> <li>(i) children aged between 1 and 3 years</li> <li>(inclusive) provides 65 mg or more total</li> <li>magnesium from inorganic magnesium</li> <li>salts;</li> </ul>
			<ul> <li>(ii) children aged between 4 and 8 years</li> <li>(inclusive) provides 110 mg or more</li> <li>total magnesium from inorganic</li> <li>magnesium salts; or</li> </ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is required on the medicine label:
			- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is oral, the medicine must not be directed for us in infants younger than 12 months of age.
3148	MAGNESIUM CHLORIDE HEXAHYDRATE	A, E, H	Magnesium is a mandatory component of magnesium chloride hexahydrate. When used in a medicine:
			(a) with an oral route of administration;
			<ul><li>(a) which an oral route of administration,</li><li>(b) not indicated for laxative (or related)</li><li>use; and</li></ul>

Volume	4

			(c) where the maximum recommended daily dose for:
			(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;
			<ul><li>(ii) children aged between 4 and 8 years</li><li>(inclusive) provides 110 mg or more</li><li>total magnesium from inorganic</li><li>magnesium salts; or</li></ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is
			required on the medicine label: - (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.
3149	MAGNESIUM CITRATE	A, E, H	
3150	MAGNESIUM CITRATE NONAHYDRATE	A, E, H	
3151	MAGNESIUM CITRATE TETRADECAHYDRATE	A, E, H	
3152	MAGNESIUM DIGLUTAMATE	A, E, H	
3153	MAGNESIUM GLUCONATE	A, E, H	
3154	MAGNESIUM GLYCEROPHOSPHATE	A, E, H	
3155	MAGNESIUM GLYCINATE	A	Only for use in oral medicines.
3156	MAGNESIUM GLYCINATE DIHYDRATE	A	Only for use in oral medicines.Magnesium is a mandatory component of Magnesium glycinate dihydrate.The percentage of Magnesium from Magnesium glycinate dihydrate should be calculated based on the molecular weight of Magnesium glycinate dihydrate.

Therapeutic Goods (Permissible Ingredients) Determination (No. 3) 2024

3157	MAGNESIUM HYDROGEN PHOSPHATE	Н	Magnesium is a mandatory component of magnesium hydrogen phosphate.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			<ul> <li>(i) children aged between 1 and 3 years</li> <li>(inclusive) provides 65 mg or more total</li> <li>magnesium from inorganic magnesium</li> <li>salts;</li> </ul>
			<ul><li>(ii) children aged between 4 and 8 years</li><li>(inclusive) provides 110 mg or more</li><li>total magnesium from inorganic</li><li>magnesium salts; or</li></ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is required on the medicine label:
			- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.
3158	MAGNESIUM HYDROXIDE	A, E	When used as an active ingredient, can only be supplied as an uncompounded medicine substance packed for retail sale, and must comply with an uncompounded substance monograph of the British Pharmacopoeia, as in force or existing from time to time.
			Magnesium is a mandatory component of magnesium hydroxide.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and

Volume 4

Vol	ume	4

			(c) where the maximum recommended daily dose for:
			<ul><li>(i) children aged between 1 and 3 years</li><li>(inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;</li></ul>
			<ul><li>(ii) children aged between 4 and 8 years</li><li>(inclusive) provides 110 mg or more</li><li>total magnesium from inorganic</li><li>magnesium salts; or</li></ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is required on the medicine label:
			- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.
3159	MAGNESIUM LYSINATE	A	Only for use in oral medicines.
3160	MAGNESIUM METHIONINATE	A	Only for use in oral medicines.
3161	MAGNESIUM NITRATE	Е	Only for use in topical medicines for dermal application.
3162	MAGNESIUM OROTATE	A, E, H	
3163	MAGNESIUM OROTATE DIHYDRATE	A, E, H	
3164	MAGNESIUM OXIDE	A, E, H	Magnesium is a mandatory component of magnesium oxide.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:

			<ul> <li>magnesium from inorganic magnesium salts;</li> <li>(ii) children aged between 4 and 8 years</li> <li>(inclusive) provides 110 mg or more total magnesium from inorganic magnesium salts; or</li> <li>(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;</li> <li>the following warning statement is required on the medicine label:</li> <li>(LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> <li>When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.</li> </ul>
3165	MAGNESIUM PHOSPHATE PENTAHYDRATE	A, E, H	<ul> <li>Magnesium is a mandatory component of magnesium phosphate pentahydrate.</li> <li>When used in a medicine: <ul> <li>(a) with an oral route of administration;</li> <li>(b) not indicated for laxative (or related) use; and</li> <li>(c) where the maximum recommended daily dose for:</li> <li>(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total magnesium from inorganic magnesium</li> </ul> </li> </ul>
			salts; (ii) children aged between 4 and 8 years (inclusive) provides 110 mg or more total magnesium from inorganic magnesium salts; or (iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;

Volume 4

(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total

Volume 4

			<ul> <li>the following warning statement is required on the medicine label:</li> <li>(LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> <li>When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.</li> </ul>
3166	MAGNESIUM PHOSPHATE TRIBASIC	A, E, H	Magnesium is a mandatory component of magnesium phosphate tribasic.
			The percentage of magnesium from magnesium phosphate tribasic should be calculated based on the molecular weigh of magnesium phosphate tribasic.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;
			(ii) children aged between 4 and 8 years (inclusive) provides 110 mg or more total magnesium from inorganic magnesium salts; or
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium
			salts; the following warning statement is required on the medicine label:
			<ul> <li>- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> </ul>
			When the route of administration is oral, the medicine must not be directed for us in infants younger than 12 months of age.

3167	MAGNESIUM PYRUVATE	А	Only for use in oral medicines.
			The maximum recommended daily dose must be no more than 7 grams.
3168	MAGNESIUM STEARATE	E	
3169	MAGNESIUM SULFATE DIHYDRATE	A, E, H	When used internally, the maximum recommended daily dose must not be more than 1.5g.
			Magnesium is a mandatory component of magnesium sulfate dihydrate.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			<ul> <li>(i) children aged between 1 and 3 years</li> <li>(inclusive) provides 65 mg or more tota</li> <li>magnesium from inorganic magnesium</li> <li>salts;</li> </ul>
			<ul> <li>(ii) children aged between 4 and 8 years</li> <li>(inclusive) provides 110 mg or more</li> <li>total magnesium from inorganic</li> <li>magnesium salts; or</li> </ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is required on the medicine label:
			- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is or a the medicine must not be directed for us in infants younger than 12 months of age.
3170	MAGNESIUM SULFATE HEPTAHYDRATE	A, E, H	When used internally, the maximum recommended daily dose must not be more than 1.5 g.

Volume	4
--------	---

			Magnesium is a mandatory component of magnesium sulfate heptahydrate.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			<ul> <li>(i) children aged between 1 and 3 years</li> <li>(inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;</li> </ul>
			<ul><li>(ii) children aged between 4 and 8 years</li><li>(inclusive) provides 110 mg or more</li><li>total magnesium from inorganic</li><li>magnesium salts; or</li></ul>
			(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;
			the following warning statement is required on the medicine label:
			- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).
			When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.
3171	MAGNESIUM SULFATE MONOHYDRATE	A, E, H	When used internally, the maximum recommended daily dose must not be more than 1.5 g.
			Magnesium is a mandatory component of magnesium sulfate monohydrate.
			When used in a medicine:
			(a) with an oral route of administration;
			(b) not indicated for laxative (or related) use; and
			(c) where the maximum recommended daily dose for:
			(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total

			Volume 4
			<ul> <li>magnesium from inorganic magnesium salts;</li> <li>(ii) children aged between 4 and 8 years (inclusive) provides 110 mg or more total magnesium from inorganic magnesium salts; or</li> <li>(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;</li> <li>the following warning statement is required on the medicine label: <ul> <li>(LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> </ul> </li> <li>When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.</li> </ul>
3172	MAGNESIUM SULFATE TRIHYDRATE	A, E, H	<ul> <li>When used internally, the maximum recommended daily dose must not be more than 1.5 g.</li> <li>Magnesium is a mandatory component of magnesium sulfate trihydrate.</li> <li>When used in a medicine: <ul> <li>(a) with an oral route of administration;</li> <li>(b) not indicated for laxative (or related) use; and</li> <li>(c) where the maximum recommended daily dose for:</li> <li>(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;</li> <li>(ii) children aged between 4 and 8 years (inclusive) provides 110 mg or more total magnesium salts; or</li> <li>(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;</li> </ul> </li> </ul>

Vol	lume	4
	CHILLE	

			<ul> <li>the following warning statement is required on the medicine label:</li> <li>- (LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> <li>When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.</li> </ul>
3173	MAGNESIUM TRISILICATE	E	<ul> <li>Magnesium is a mandatory component of magnesium trisilicate.</li> <li>When used in a medicine: <ul> <li>(a) with an oral route of administration;</li> <li>(b) not indicated for laxative (or related) use; and</li> <li>(c) where the maximum recommended daily dose for:</li> <li>(i) children aged between 1 and 3 years (inclusive) provides 65 mg or more total magnesium from inorganic magnesium salts;</li> <li>(ii) children aged between 4 and 8 years (inclusive) provides 110 mg or more total magnesium salts; or</li> <li>(iii) individuals aged 9 years or older provides 350 mg or more total magnesium from inorganic magnesium salts;</li> <li>the following warning statement is required on the medicine label: <ul> <li>(LAX6) 'Contains magnesium, which may have a laxative effect or cause diarrhoea' (or words to that effect).</li> </ul> </li> <li>When the route of administration is oral, the medicine must not be directed for use in infants younger than 12 months of age.</li> </ul></li></ul>
3174	MAGNOLIA GLAUCA	A, H	
3175	MAGNOLIA LILIFLORA	A, H	
5115	MAUNULIA LILIFLUKA	1,11	

## Volume 4

3177	MAGNOLIA OFFICINALIS	A, E, H	
3178	MAGNOLIA SALICIFOLIA	A, H	
3179	MAIZE OIL	A, E, H	
3180	MAIZE STARCH	A, E, H	
3181	MALACHITE GREEN	Е	Permitted for use only as a colour for topical use.
3182	MALIC ACID	E	Sponsors should consider the impact of excipients on the sensitivity of the skin to sunlight and should ensure the finished medicine is safe for its intended purpose.
3183	MALPIGHIA GLABRA	A, E, H	
3184	MALT EXTRACT	Е	
3185	MALTITOL	Е	
3186	MALTITOL SOLUTION	Е	
3187	MALTODEXTRIN	E	Gluten is a mandatory component of Maltodextrin where the ingredient is derived from gluten containing grains such as wheat, barley, rye and oats.
3188	MALTOL	E	
3189	MALTONE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3190	MALTOSE	E	
3191	MALUS DOMESTICA	A, E, H	The concentration of amygdalin in the medicine must be no more than 0%.
3192	MALUS SYLVESTRIS	A, H	
3193	MALVA MOSCHATA	A, H	
3194	MALVA SYLVESTRIS	A, E, H	
3195	MALVA VERTICILLATA	A, H	
3196	MANDARIN	Е	

Volun	ne	4

3197	MANDARIN OIL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3198	MANDARIN OIL COLDPRESSED	A, E, H	When used internally, oxedrine is a mandatory component of mandarin oil coldpressed.
			The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
3199	MANDARIN OIL TERPENES	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3200	MANDARIN RESIDUE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3201	MANDRAGORA OFFICINARUM	A, H	Atropine, hyoscine and hyoscyamine are mandatory components of Mandragora officinarum.
			The concentration in the medicine must be no more than 10 mg/kg or 10 mL/L or 0.001%.
			The concentration of atropine in the medicine must be no more than 100

			Volume 4
			micrograms/kg or 100 micrograms/L or 0.00001%.
			The concentration of hyoscine in the medicine must be no more than 300 micrograms/kg or 300 micrograms/L or 0.00003%.
			The concentration of hyoscyamine in the medicine must be no more than 300 micrograms/kg or 300 micrograms/L or 0.00003%.
3202	MANGANESE	Н	Only for use as an active homoeopathic ingredient.
3203	MANGANESE (II) DIASPARTATE	A, H	Only for use in oral medicines.
3204	MANGANESE (II) GLYCINATE	A, H	Only for use in oral medicines.
3205	MANGANESE ACETATE TETRAHYDRATE	Н	Only for use as an active homoeopathic ingredient.
3206	MANGANESE AMINO ACID CHELATE	A, E, H	Only for use in oral medicines. The concentration of Manganese must be no more than 25% of the manganese amino acid chelate.
3207	MANGANESE CHLORIDE TETRAHYDRATE	A, E, H	
3208	MANGANESE DIASPARTATE	A, E, H	Only for use in oral medicines.
3209	MANGANESE GLUCONATE	A, E, H	
3210	MANGANESE GLYCEROPHOSPHATE	A, E, H	
3211	MANGANESE OXIDE	A, E, H	
3212	MANGANESE SULFATE MONOHYDRATE	A, E, H	
3213	MANGANESE SULFATE TETRAHYDRATE	A, E, H	
3214	MANGIFERA INDICA	A, E, H	
3215	MANGO	E, H	
3216	MANIHOT ESCULENTA	A, H	
3217	MANNITOL	Е	

Volume 4	ŀ
----------	---

3218	MARANTA ARUNDINACEA	A, H	
3219	MARINE SPONGE	Н	Only for use as an active homoeopathic ingredient.
3220	MARJORAM OIL SPANISH	A, E, H	<ul> <li>When the concentration in the preparation is more than 50%, the nominal capacity of the container must be no more than 50 mL, the medicine must have a restricted flow insert fitted to the container and requires the following warning statement on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect).</li> </ul>
3221	MARJORAM OIL SWEET	A, E, H	<ul> <li>When the concentration in the preparation is more than 50%, the nominal capacity of the container must be no more than 50 mL, the medicine must have a restricted flow insert fitted to the container and requires the following warning statement on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect).</li> </ul>
3222	MARRUBIUM VULGARE	A, E, H	
3223	MARSDENIA CUNDURANGO	A, H	
3224	MARSHMALLOW ROOT DRY	A, H	
3225	MARSHMALLOW ROOT POWDER	A, H	
3226	MASSOIA LACTONE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3227	MASTIC	A, H	
3228	MATE ABSOLUTE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.

			Volume
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3229	MATRICARIA CHAMOMILLA	A, E, H	
3230	MATRICARIA FLOWER DRY	A, E, H	
3231	MEADOWSWEET HERB DRY	А, Н	Methyl salicylate is a mandatory component of meadowsweet herb dry.
			Not to be included in medicines for use in the eye or on damaged skin.
			When used internally, the concentration of methyl salicylate in the medicine must not be more than 0.001%.
		When the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is other than spray, the medicine requires child resistant packaging.	
			<ul> <li>When the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is spray, the medicine does not require child resistant packaging if:</li> </ul>
			- the delivery device is engaged into the container in such a way that prevents it from being readily removed;
		- direct suction through the delivery device results in delivery of no more that one dosage unit; and	
		- actuation of the spray device is ergonomically difficult for young children to accomplish.	
		The following warning statement is required on the medicine label:	
			- (METSAL) 'Contains methyl salicylat (or words to that effect).
			When for use in topical medicines for dermal application

Volum	le 4

			<ul><li>i) the concentration of methyl salicylate</li><li>in the medicine must not be more than</li><li>25%</li></ul>
			ii) the following warning statements are required on the medicine label:
			- (PREGNT2) 'Do not use if pregnant or likely to become pregnant' (or words to that effect);
			- (CHILD4) 'Do not use [this product/insert name of product] in children 6 years of age or less';
			- (SENS) 'Application to skin may increase sensitivity to sunlight.' (or words to that effect);
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to that effect);
			<ul><li>iii) if the concentration of methyl salicylate in the medicine is greater than 1%, the following warning statement is required on the medicine label:</li></ul>
			- (IRRIT) 'If irritation develops, discontinue use'.
3232	MECOBALAMIN	A	Only for use in oral medicines.
3233	MEDICAGO SATIVA	A, E, H	The level of l-canavanine must be no more than that of the dried leaf.
			When fresh leaf extract is used and the extraction ratio is between 34:1 and 46:1, the quantity of l-canavanine in the extract must not be more than that in the fresh leaf.
3234	MEDIUM CHAIN TRIGLYCERIDES	E	
3235	MELALEUCA ALTERNIFOLIA	A, E, H	Cineole is a mandatory component of Melaleuca alternifolia.
			In liquid preparations when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:
			a) the nominal capacity of the container must be no more than 25 millilitres;

48

			Volume 4
			<ul> <li>b) a restricted flow insert must be fitted on the container; and</li> <li>c) the container must include the following warning statements on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>- (NTAKEN) 'Not to be taken'.</li> <li>In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres the medicine must also have a child resistant closure.</li> </ul>
3236	MELALEUCA CAJUPUTI	A, E, H	<ul> <li>Cineole is a mandatory component of Melaleuca cajuputi.</li> <li>In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:</li> <li>a) the nominal capacity of the container must be no more than 25 millilitres;</li> <li>b) a restricted flow insert must be fitted on the container; and</li> <li>c) the container must include the following warning statements on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>- (NTAKEN) 'Not to be taken'.</li> <li>In liquid preparations, when the concentration of cineole OR the concentration of cineole OR the concentration of cillor distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres the medicine must also have a child resistant closure.</li> </ul>
3237	MELALEUCA CITRINA	A, H	

Volume	4
--------	---

3238	MELALEUCA DISSITIFLORA	А, Н	Cineole is a mandatory component of Melaleuca dissitiflora.
			In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:
			a) the nominal capacity of the container must be no more than 25 millilitres;
			b) a restricted flow insert must be fitted on the container; and
			c) the container must include the following warning statements on the medicine label:
			<ul> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>- (NTAKEN) 'Not to be taken'.</li> </ul>
			In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres the medicine must also have a child resistant closure.
3239	MELALEUCA ERICIFOLIA	A, E, H	Cineole is a mandatory component of Melaleuca ericifolia.
			In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:
			a) the nominal capacity of the container must be no more than 25 millilitres;
			b) a restricted flow insert must be fitted on the container; and
			c) the container must include the following warning statements on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect); and
			- (NTAKEN) 'Not to be taken'.
			In liquid preparations, when the concentration of cineole OR the

Volume 4

			Volume 4
			concentration of oil or distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres the medicine must also have a child resistant closure.
3240	MELALEUCA LINARIIFOLIA	A, H	<ul> <li>Cineole is a mandatory component of Melaleuca linariifolia.</li> <li>In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:</li> <li>a) the nominal capacity of the container must be no more than 25 millilitres;</li> <li>b) a restricted flow insert must be fitted on the container; and</li> <li>c) the container must include the following warning statements on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>- (NTAKEN) 'Not to be taken'.</li> <li>In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres the medicine must also have a child resistant closure.</li> </ul>
3241	MELALEUCA OIL	A, E, H	Cineole and cajuput oil are a mandatory components of Melaleuca Oil. When the plant preparation is oil and the concentration in the medicine is more than 25%, the nominal capacity of the container must be no more than 25 mL and the medicine requires the following warning statements on the medicine label: - (CHILD) 'Keep out of reach of children' (or word to that effect) - (NTAKEN) 'Not to be taken'.

Volume 4

			When the nominal capacity of the container is 15 mL or less, then a restricted flow insert must be fitted on the container.
			Where the nominal capacity of the container is more than 15 mL but less than or equal to 25 mL, then a child resistant closure and restricted flow insert must be fitted on the container.
3242	MELALEUCA QUINQUENERVIA	A, E, H	Cineole is a mandatory component of Melaleuca quinquenervia.
			In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25%:
			a) the nominal capacity of the container must be no more than 25 millilitres;
			b) a restricted flow insert must be fitted on the container; and
			c) the container must include the following warning statements on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect); and
			- (NTAKEN) 'Not to be taken'.
			In liquid preparations, when the concentration of cineole OR the concentration of oil or distillate in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres the medicine must also have a child resistant closure.
3243	MELICOPE PTELEIFOLIA	A, H	
3244	MELILOTUS OFFICINALIS	A, E, H	Coumarin is a mandatory component of Melilotus officinalis.
			The concentration of coumarin in the medicine must be no more than 0.001%.
3245	MELISSA OFFICINALIS	A, E, H	
3246	MELON	Е	

Vol	ume	4
<b>V</b> U	unit	т

3247	MENADIONE SODIUM BISULFITE	Е	
3248	MENAQUINONE 7	A	For oral use only. The medicine must not provide more than 180 micrograms per maximum daily dose in adults, 90 micrograms per maximum daily dose in children between 10-18 years, and 45 micrograms per maximum daily dose in children less than 10 years of age.
3249	MENISPERMUM CANADENSE	A, H	
3250	MENTHA AQUATICA	A, H A, H	<ul> <li>Menthol is a mandatory component of Mentha aquatica.</li> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statement is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> </ul> </li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the</li> </ul></li></ul>
			<ul> <li>following warning statements are required on the medicine label:</li> <li>- (SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>- (IRRIT) If irritation develops, discontinue use.</li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label:</li> <li>- (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> </ul>

Volume 4	4
----------	---

			When the medicine is for internal use, the maximum recommended daily dose must not contain more than 1 gram of menthol.
3251	MENTHA ARVENSIS	A, E, H	<ul> <li>Menthol is a mandatory component of Mentha arvensis.</li> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statement is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label: <ul> <li>(IRRIT) If irritation develops, discontinue use.</li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label: <ul> <li>(MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> </ul> </li> </ul></li></ul></li></ul></li></ul>
3252	MENTHA ARVENSIS LEAF OIL	E	Permitted for use only in combination with other permitted ingredients as a

	flavour proprietary excipient formul or fragrance proprietary excipient formulation.	ation
	The total flavour proprietary excipie formulation in a medicine must be n more than 5%.	
	The total fragrance proprietary excip formulation in a medicine must be n more 1%.	-
	Menthol is a mandatory component Mentha arvensis leaf oil.	of
	When the medicine is for topical use dermal application:	e for
	(i) the medicine must not be intended use in the eye or on damaged skin;	d for
	(ii) the medicine must not deliver me than 25% total menthol when administered according to the direct for use;	
	(iii) the following warning statement required on the medicine label:	t is
	- (EYE) Avoid contact with eyes (or words to that effect).	
	<ul> <li>(iv) if the medicine delivers more that</li> <li>1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label:</li> </ul>	d
	- (SKTEST) If you have sensitive sk test this product on a small area of sl before applying it to a large area;	
	- (IRRIT) If irritation develops, discontinue use.	
	(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is require on the medicine label:	d :he
	- (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.	1
	When the medicine is for internal us the maximum recommended daily d	-

## Volume 4

			must not contain more than 1 gram of menthol.
3253	MENTHA ARVENSIS OIL	E	<ul> <li>menthol.</li> <li>Permitted for use only in combination with other permitted ingredients as a flavour proprietary excipient formulation.</li> <li>The total flavour proprietary excipient formulation in a medicine must not be more than 5%.</li> <li>Menthol is a mandatory component of Mentha arvensis oil.</li> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statements is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> </ul> </li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> </ul> </li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> </ul></li></ul>
			following warning statement is required on the medicine label: - (MENTH) Contains a high
			concentration of menthol, which can cause severe skin irritation.
			When the medicine is for internal use, the maximum recommended daily dose

			must not contain more than 1 gram of menthol.
3254	MENTHA HAPLOCALYX	A, E, H	Menthol is a mandatory component of Mentha haplocalyx.
			When the medicine is for topical use for dermal application:
			(i) the medicine must not be intended fo use in the eye or on damaged skin;
			<ul><li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li></ul>
			(iii) the following warning statement is required on the medicine label:
			- (EYE) Avoid contact with eyes (or words to that effect).
			(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label:
			- (SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;
			- (IRRIT) If irritation develops, discontinue use.
			<ul> <li>(v) if the medicine delivers more than</li> <li>5% total menthol when administered</li> <li>according to the directions for use, the</li> <li>following warning statement is required</li> <li>on the medicine label:</li> </ul>
			- (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.
			When the medicine is for internal use, the maximum recommended daily dose must not contain more than 1 gram of menthol.
3255	MENTHA PULEGIUM	А, Н	D-pulegone, menthol and volatile oil components (of Mentha pulegium) are mandatory components of Mentha pulegium.

Volume	4
--------	---

<ul> <li>When the nominal capacity of the container is more than 15 millilitres, the concentration of d-pulegone in the medicine must be no more than 4%.</li> <li>When the concentration of d-pulegone in the preparation is more than 4% and the nominal capacity of the container is 15 millilitres or less, the medicine must have a child resistant closure and restricted flow insert fitted on the container.</li> <li>The medicine requires the following warning statements on the medicine label: <ul> <li>(NTAKEN) 'Not to be taken';</li> <li>(CHILD) 'Keep out of reach of children' (or words to that effect).</li> </ul> </li> <li>When the medicine is for topical use for dermal application: <ul> <li>a) the maximum recommended daily dose must not contain more than 150 mg</li> </ul> </li> </ul>
restricted flow insert fitted on the container. The medicine requires the following warning statements on the medicine label: - (NTAKEN) 'Not to be taken'; - (CHILD) 'Keep out of reach of children' (or words to that effect). When the medicine is for topical use for dermal application:
of Mentha pulegium oil or distillate; b) the medicine must not be intended for use in the eye or on damaged skin;
c) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;
d) the following warning statement is required on the medicine label:
- (EYE) Avoid contact with eyes (or words to that effect).
e) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label:
- (SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;
- (IRRIT) If irritation develops, discontinue use.
f) if the medicine delivers more than 5% total menthol when administered

59

Therapeutic Goods (Permissible Ingredients) Determination (No. 3) 2024

			<ul> <li>according to the directions for use, the following warning statement is required on the medicine label:</li> <li>– (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> <li>When the medicine is for internal use:</li> <li>a) the maximum recommended daily dose must not contain more than 50 mg of Mentha pulegium oil or distillate;</li> <li>b) the maximum recommended daily dose must not contain more than 1 gram of menthol.</li> </ul>
3256	MENTHA SPICATA	A, E, H	<ul> <li>Menthol is a mandatory component of Mentha spicata.</li> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statement is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> </ul> </li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine delivers more than</li> </ul></li></ul>

Volume 4

Volume 4
----------

			<ul> <li>(MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> <li>When the medicine is for internal use, the maximum recommended daily dose must not contain more than 1 gram of menthol.</li> </ul>
3257	MENTHA X CARDIACA	A, E, H	<ul> <li>Menthol is a mandatory component of Mentha x cardiaca.</li> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statement is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label: <ul> <li>(IRRIT) If on the irritation develops, discontinue use.</li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label: <ul> <li>(MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> </ul> </li> </ul></li></ul></li></ul></li></ul>
			<ul> <li>following warning statement is require</li> <li>on the medicine label:</li> <li>– (MENTH) Contains a high</li> <li>concentration of menthol, which can</li> </ul>

3258	MENTHA X PIPERITA	A, E, H	Menthol is a mandatory component of Mentha x piperita.
			When the medicine is for topical use for dermal application:
			(i) the medicine must not be intended fo use in the eye or on damaged skin;
			<ul><li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li></ul>
			(iii) the following warning statement is required on the medicine label:
			- (EYE) Avoid contact with eyes (or words to that effect).
			<ul> <li>(iv) if the medicine delivers more than</li> <li>1% total menthol when administered</li> <li>according to the directions for use, the</li> <li>following warning statements are</li> <li>required on the medicine label:</li> </ul>
			- (SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;
			- (IRRIT) If irritation develops, discontinue use.
			<ul> <li>(v) if the medicine delivers more than</li> <li>5% total menthol when administered</li> <li>according to the directions for use, the</li> <li>following warning statement is required</li> <li>on the medicine label:</li> </ul>
			– (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.
			When the medicine is for internal use, the maximum recommended daily dose must not contain more than 1 gram of menthol.
3259	MENTHADIENYL ACETATE	E	Menthadienyl acetate must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.

Volume 4

			The total concentration of the flavour proprietary excipient formulation containing menthadienyl acetate must not be more than 5% of the total medicine.
3260	MENTHANYL ACETATE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3261	MENTHOFURAN	Е	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3262	MENTHOL	A, E	<ul> <li>When the medicine is for topical use for dermal application: <ul> <li>(i) the medicine must not be intended for use in the eye or on damaged skin;</li> <li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li> <li>(iii) the following warning statement is required on the medicine label: <ul> <li>(EYE) Avoid contact with eyes (or words to that effect).</li> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> </ul></li></ul></li></ul>

			<ul> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label:</li> <li>– (MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> <li>When the medicine is for internal use, the maximum recommended daily dose must not contain more than 1 gram of menthol.</li> </ul>
3263	MENTHONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3264	MENTHONE GLYCERINE ACETAL	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3265	MENTHONE THIOL FRACTION	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3266	MENTHOXYPROPANEDIOL	Е	For oral use only. The concentration in the medicine must be no more than 0.04%.

Volume	4

3267	MENTHYL 2-HYDROXYETHYL CARBONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3268	MENTHYL 2-HYDROXYPROPYL CARBONATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3269	MENTHYL ANTHRANILATE	A	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must not be more than 5%.
			<ul> <li>When used in primary sunscreen products, the following warning statements are required on the label:</li> <li>- (AVOID) 'Avoid prolonged exposure in the sun' (or words to this effect); and</li> <li>- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).</li> </ul>
3270	MENTHYL ISOVALERATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3271	MENTHYL LACTATE	E	
3272	MENYANTHES TRIFOLIATA	A, H	
3273	MERCURIC CHLORIDE	Н	Only for use as an active homoeopathic ingredient.
			Mercury is a mandatory component of mercuric chloride.

			The total concentration of mercury in the medicine must not be more than 1 mg/kg or 1 mg/L or 0.0001%.
3274	MERCURY	H	Only for use as an active homoeopathic ingredient. The total concentration of mercury in the medicine must not be more than 1 mg/kg or 1 mg/L or 0.0001%.
3275	METACRESOL	E	Only for use in topical medicines for dermal application.
3276	METHACRYLIC ACID COPOLYMER	E	Only for use in oral medicines.
3277	METHANOL	E	The residual solvent limit is 30 mg per recommended daily dose. The concentration in the medicine must be no more than 0.3%.
3278	METHICONE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 1%.
3279	METHIONINE	A, E	
3280	METHYL 2,6,6- TRIMETHYLCYCLOHEX-2-ENE-1- CARBOXYLATE	E	<ul> <li>Only for use in medicines in combination with other permitted ingredients as a fragrance proprietary excipient formulation.</li> <li>The total fragrance proprietary excipient formulation in a medicine must not be more than 1%.</li> </ul>
3281	METHYL 2-METHYLBUTYRATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>

Volume 4

			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3282	METHYL 2-OCTYNOATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3283	METHYL 3,6- DIMETHYLRESORCYLATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3284	METHYL ACETATE	E	The residual solvent limit is 50 mg per recommended daily dose.
			The concentration in the medicine must be no more than 0.5%.
3285	METHYL ACETOPHENONE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3286	METHYL ACETYL RICINOLEATE	E	Only for use in topical medicines for dermal application.
3287	METHYL ANISATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.

			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3288	METHYL ANTHRANILATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3289	METHYL BENZOATE	E	Only for use in topical medicines for dermal application.
3290	METHYL BUTYRATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3291	METHYL CAPROATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3292	METHYL CAPRYLATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>

Volume	4

3293	METHYL CARBITOL	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance</li></ul>
			concentration in a medicine must be no more than 1%.
3294	METHYL CEDRYL KETONE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3295	METHYL CHAVICOL	E	Permitted for use only in combination with other permitted ingredients as part of a fragrance proprietary excipient formulation.
			The ingredient is not to be included in medicines intended for oral use.
			The quantity of methyl chavicol in a medicine must be no more than 0.01%.
			The total fragrance proprietary excipien formulation in a medicine must be no more than 1%.
3296	METHYL CINNAMATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3297	METHYL CIS-5-OCTENOATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

2208	METHVI CVCI OPENITENOI ONE	Б	Domitted for use only in combination
3298	METHYL CYCLOPENTENOLONE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3299	METHYL CYCLOPENTYLIDENEACETATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3300	METHYL DI-TERT-BUTYL-4- HYDROXYHYDROCINNAMATE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3301	METHYL DIHYDROABIETATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3302	METHYL DIISOPROPYL PROPIONAMIDE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3303	METHYL ETHER	E	Only for use in topical medicines for dermal application.
3304	METHYL ETHYL KETONE	E	The residual solvent limit is 50 mg per maximum recommended daily dose.

Volume -	4
----------	---

			The concentration in the medicine must be no more than 0.5%.
3305	METHYL EUGENOL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3306	METHYL FUROATE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3307	METHYL GLUCETH-10	E	<ul> <li>Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.</li> <li>The concentration in the medicine must be no more than 3%.</li> <li>Residue levels of ethylene oxide are to be kept below the level of detection.</li> </ul>
3308	METHYL GLUCETH-20	E	Only for use in topical medicines for dermal application.
3309	METHYL GLUCETH-20 BENZOATE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3310	METHYL GLUCETH-20 SESQUIHYDRATE	E	Only for use in topical medicines for dermal application.

3311	METHYL GLUCOSE DIOLEATE	E	Only for use in topical medicines for dermal application.
3312	METHYL GLUCOSE SESQUIOLEATE	E	Only for use in topical medicines for dermal application.
3313	METHYL GLUCOSE SESQUISTEARATE	E	Only for use in topical medicines for dermal application.
3314	METHYL HEPTANOATE	E	Permitted for use only in combination with other permitted ingredients as a flavour proprietary excipient formulation.The total flavour proprietary excipient 
3315	METHYL HEPTENONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3316	METHYL HEPTYL KETONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3317	METHYL HEXYL CARBINOL	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

## Volume 4

3318	METHYL HEXYL KETONE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3319	METHYL HYDROGENATED ROSINATE	E	Only for use in topical medicines for dermal application.
3320	METHYL HYDROJASMONATE	E	Only for use in topical medicines for dermal application.
3321	METHYL HYDROXYBENZOATE	E	
3322	METHYL IONONE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3323	METHYL ISOBUTYL KETONE	E	The residual solvent limit is 50 mg per maximum daily dose.
			The concentration in the medicine must be no more than 0.5%.
3324	METHYL ISOEUGENOL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.

3325	METHYL ISOVALERATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3326	METHYL JASMONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3327	METHYL LAURATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used as a flavour the total flavour concentration in a medicine must be no more than 5%.
3328	METHYL LINOLEATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3329	METHYL LINOLENATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3330	METHYL MAGNESIUM CHLORIDE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume -	4
----------	---

3331	METHYL METHACRYLATE CROSSPOLYMER	E	Methyl methacrylate is a mandatory component of methyl methacrylate crosspolymer.
			Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			When the concentration of methyl methacrylate crosspolymer is greater than 1%, the medicine must not be
			intended for use on damaged skin. The total concentration of methyl methacrylate crosspolymer in the medicine must not be more than 4.85%.
			The total concentration of methyl methacrylate as residual monomer in the medicine must not be more than 1%.
3332	METHYL METHOXY PYRAZINE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3333	METHYL MYRISTATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3334	METHYL NAPHTHYL KETONE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3335	METHYL NONYL KETONE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.

			If used in a flavour the total flavour concentration in a medicine must be no more than 5%. If used in a fragrance the total fragrance
			concentration in a medicine must be no more 1%.
3336	METHYL NONYLENATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3337	METHYL OCTIN CARBONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3338	METHYL PALMITATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3339	METHYL PHENYL CARBINOL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume	4

3340	METHYL PHENYL CARBINYL-ISO- BUTYRATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no</li></ul>
3341	METHYL PHENYL GLYCIDATE	E	more than 5%. Permitted for use only in combination
			<ul><li>with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
3342	METHYL PHENYLACETATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3343	METHYL PHENYLCARBINYL ACETATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3344	METHYL ROSINATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3345	METHYL SALICYLATE	A, E	Not to be included in medicines for use in the eye or on damaged skin.

Vo	lume	4
VU.	unit	т.

When used internally, the concentration in the medicine must not be more than 0.001%.
When the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is other than spray, the medicine requires child resistant packaging.
When the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is spray, the medicine does not require child resistant packaging if:
- the delivery device is engaged into the container in such a way that prevents it from being readily removed;
- direct suction through the delivery device results in delivery of no more than one dosage unit; and
- actuation of the spray device is ergonomically difficult for young children to accomplish.
The following warning statement is required on the medicine label:
- (METSAL) 'Contains methyl salicylate' (or words to that effect).
When for use in topical medicines for dermal application:
i) the concentration of methyl salicylate in the medicine must not be more than 25%;
ii) the following warning statements are required on the medicine label:
- (PREGNT2) 'Do not use if pregnant or likely to become pregnant' (or words to that effect);
- (CHILD4) 'Do not use [this product/insert name of product] in children 6 years of age or less';
- (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect);
- (AVOID) 'Avoid prolonged exposure in the sun' (or words to that effect);

Volume 4

			<ul> <li>iii) if the concentration of methyl salicylate in the medicine is greater than 1%, the following warning statement is required on the medicine label:</li> <li>- (IRRIT) 'If irritation develops, discontinue use'.</li> </ul>
3346	METHYL STEARATE	Е	
3347	METHYL THIOBUTYRATE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3348	METHYL TRIMETICONE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 5%.
3349	METHYL-3- METHYLTHIOPROPIONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3350	METHYL-BETA-METHYL THIOLPROPIONATE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3351	METHYL-PARA-TERT-BUTYL PHENYLACETATE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume 4

3352	METHYLBENZYL ACETATE	E	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3353	METHYLCELLULOSE	A, E	
3354	METHYLCHLOROISOTHIAZOLINON E	E	Only for use in topical medicines for dermal application that are rinsed off the skin.
			The total concentration of methylchloroisothiazolinone and methylisothiazolinone in the medicine must be no more than 0.0015%.
3355	METHYLCYCLOHEXADIENE	E	Permitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance
			concentration in a medicine must be no more than 1%.
3356	METHYLENE BIS- BENZOTRIAZOLYL TETRAMETHYLBUTYLPHENOL	A	Only for use as an active ingredient in sunscreens for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must not be more than 10%.
			When used in primary sunscreen products, the following warning statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).
3357	METHYLISOTHIAZOLINONE	E	Only for use in topical medicines for dermal application that are rinsed off the skin.
			The total concentration of methylchloroisothiazolinone and

Volume 4

			methylisothiazolinone in the medicine must be no more than 0.0015%.
3358	METHYLMERCAPTAN	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3359	METHYLPROPANEDIOL	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 10%.
3360	METHYLSILANOL/SILICATE CROSSPOLYMER	E	<ul> <li>Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or on damaged skin.</li> <li>The concentration in the medicine must be no more than 0.1%.</li> </ul>
3361	METHYLSTYRENE/VINYLTOLUENE COPOLYMER	E	Only for use in topical medicines for dermal application.
3362	MICA	E	Only for use when the route of administration is oral, dental or topical. The concentration in oral medicines must be no more than 2.5%. The concentration in dental toothpastes must be no more than 0.5%.
3363	MICROCALICIUM ARENARIUM	A, H	
3364	MICROCOCCUS LUTEUS LYSATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.005%.
3365	MICROCOS PANICULATA	A, H	
3366	MICROCRYSTALLINE CELLULOSE	E	

3367	MICROCRYSTALLINE WAX	E	Only for use as an excipient in medicines for topical, oral or oral application routes of administration.
			When microcrystalline wax is used as an excipient ingredient, the route of administration 'oral' is only permitted when the dosage form is 'chewing gum'.
3368	MILK FAT	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3369	MILK THISTLE FRUIT DRY	A, H	
3370	MILK THISTLE FRUIT POWDER	A, H	
3371	MILLET	Е	
3372	MILLETTIA DIELSIANA	A, H	
3373	MIMOSA ABSOLUTE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3374	MIMULUS GUTTATUS	A, H	
3375	MINT OIL DEMENTHOLISED	A, E, H	Menthol is a mandatory component of mint oil dementholised.
			When the medicine is for topical use for dermal application:
			(i) the medicine must not be intended for use in the eye or on damaged skin;
			<ul><li>(ii) the medicine must not deliver more than 25% total menthol when administered according to the directions for use;</li></ul>
			(iii) the following warning statement is required on the medicine label:
			- (EYE) Avoid contact with eyes (or words to that effect).

Volume	4
--------	---

3376MINTLACTONEEPermitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.3377MITCHELLA REPENSA, H3378MIXED (HIGH-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3379MIXED (LOW-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3380MIXED TERPENESEPermitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no				<ul> <li>(iv) if the medicine delivers more than 1% total menthol when administered according to the directions for use, the following warning statements are required on the medicine label: <ul> <li>(SKTEST) If you have sensitive skin, test this product on a small area of skin before applying it to a large area;</li> <li>(IRRIT) If irritation develops, discontinue use.</li> </ul> </li> <li>(v) if the medicine delivers more than 5% total menthol when administered according to the directions for use, the following warning statement is required on the medicine label: <ul> <li>(MENTH) Contains a high concentration of menthol, which can cause severe skin irritation.</li> </ul> </li> </ul>
3377MITCHELLA REPENSA, H3378MIXED (HIGH-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3379MIXED (LOW-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3380MIXED TERPENESEBernitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance	3376	MINTLACTONE	Е	with other permitted ingredients as a flavour.
3378MIXED (HIGH-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3379MIXED (LOW-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3380MIXED TERPENESEBernitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance				concentration in a medicine must be no
TOCOPHEROLS CONCENTRATEA, E3379MIXED (LOW-ALPHA TYPE) TOCOPHEROLS CONCENTRATEA, E3380MIXED TERPENESEPermitted for use only in combination with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance	3377	MITCHELLA REPENS	A, H	
TOCOPHEROLS CONCENTRATE       E       Permitted for use only in combination with other permitted ingredients as a fragrance.         3380       MIXED TERPENES       E       Permitted for use only in combination with other permitted ingredients as a fragrance.	3378	· · · · · · · · · · · · · · · · · · ·	A, E	
with other permitted ingredients as a fragrance. If used in a fragrance the total fragrance	3379	· · · · · · · · · · · · · · · · · · ·	A, E	
	3380	MIXED TERPENES	E	<ul><li>with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance</li></ul>

3382	MOLASSES	Е	Permitted for use only in combination with other permitted ingredients as a
			flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3383	MOLYBDENUM	H	Only for use as an active homoeopathic ingredient. When Molybdenum is sourced from Molybdenum trioxide then the maximum daily dose must be no more than 125 micrograms. When Molybdenum is sourced from yeast - high molybdenum then the maximum recommended daily dose must be no more than 62.5 micrograms.
3384	MOLYBDENUM TRIOXIDE	A	Molybdenum is a mandatory component of Molybdenum trioxide. The maximum daily dose of molybdenum from Molybdenum trioxide must be no more than 125 micrograms. The percentage of molybdenum from molybdenum trioxide should be calculated based on the molecular weigh of molybdenum trioxide.
3385	MOMORDICA BALSAMINA	A, H	
3386	MOMORDICA CHARANTIA	A, H	
3387	MOMORDICA COCHINCHINENSIS	A, H	
3388	MONARDA DIDYMA	A, H	
3389	MONO- AND DI- GLYCERIDES	Е	
3390	MONOBASIC AMMONIUM PHOSPHATE	E	Only for use in topical medicines for dermal application.
3391	MONOBASIC CALCIUM PHOSPHATE	A, E, H	
3392	MONOBASIC POTASSIUM PHOSPHATE	A, E, H	When used in a solid medicine containing this ingredient, the pH of a 1 g/L aqueous solution must not be more than 11.5.

Volume 4

			When used in a liquid or a semi-solid medicine containing this ingredient, the pH of the medicine must be no more than 11.5.
3393	MONOBASIC SODIUM PHOSPHATE	A, E, H	When used in a solid preparation, the pH of a 10 g/L aqueous solution must not be more than 11.5.
			When used in a liquid or a semi-solid preparation, the pH of the preparation must not exceed 11.5.
3394	MONOBASIC SODIUM PHOSPHATE DIHYDRATE	E	When used in a solid preparation, the pH of a 10 g/L aqueous solution must not be more than 11.5.
			When used in a liquid or a semi-solid preparation, the pH of the preparation must not exceed 11.5.
3395	MONOETHANOLAMINE	E	Only for use in topical medicines for dermal application.
			The concentration in the medicine must be no more than 5%.
3396	MONOMENTHYL GLUTARATE	E	Monomenthyl glutarate must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.
			The total concentration of flavour proprietary excipient formulations containing monomenthyl glutarate must not be more than 5% of the total medicine.
			The maximum recommended daily dose of the medicine must not provide more than 1.8mg of monomenthyl glutarate.
3397	MONOMENTHYL SUCCINATE	E	Monomenthyl succinate must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.

			The total concentration of the flavour proprietary excipient formulation containing monomenthyl succinate must not be more than 5% of the total medicine.
3398	MONOPHOSPHOTHIAMINE	A	
3399	MONOPHOSPHOTHIAMINE DIHYDRATE	А	
3400	MONOPOTASSIUM GLUTAMATE	Α, Ε	
3401	MONOSODIUM DIHYDROGEN CITRATE	E	
3402	MONOSODIUM GLUTAMATE MONOHYDRATE	A, E	
3403	MONSTERA DELICIOSA	A, H	
3404	MONTAN WAX	Е	
3405	MORDANT RED 11	E	Permitted for use only as a colour for topical use.
			The concentration in the medicine must be no more than 0.05%
3406	MORINDA CITRIFOLIA	А, Н	Only for use when the plant part is fruit and the plant preparation is fruit juice or fruit powder. Fruit powder must be produced by freeze drying the whole fruit (excluding the seeds).
3407	MORINDA OFFICINALIS	A, H	
3408	MORINGA OLEIFERA	A, H	
3409	MORUS ALBA	A, H	
3410	MORUS BOMBYCIS	A, H	
3411	MORUS NIGRA	A, E, H	
3412	MOSKENE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
3413	MOTHERWORT HERB DRY	A, H	

Volume 4

Volume	4
--------	---

3414	MOTHERWORT HERB POWDER	A, H	
3415	MUCUNA PRURIENS	A	Levodopa is a mandatory component of Mucuna pruriens. The concentration of levodopa in the medicine must not be more than 10
			mg/kg or 10 mg/L or 0.001%.
3416	MULBERRY	Е	
3417	MUNG BEAN	Е	
3418	MURRAYA KOENIGII	A, H	
3419	MURRAYA PANICULATA	A, H	
3420	MUSA X PARADISIACA	A, H	
3421	MUSK KETONE	E	Only for use in topical medicines for dermal application.
3422	MUSK TIBETENE	Е	Permitted for use only in combination with other permitted ingredients as a fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3423	MUSK XYLOL	E	Only for use in topical medicines for dermal application.
3424	MUSKS	Н	Only for use as an active homoeopathic ingredient.
3425	MUSTARD	E	Allyl isothiocyanate is a mandatory component of mustard when the plant part is seed.
			The concentration of allyl isothiocyanat from all ingredients in the product must be no more than 10 mg/kg or 10 mg/L o 0.001%.
3426	MUSTARD OIL	E	Allyl isothiocyanate is a mandatory component of mustard oil when the plan part is seed.
			The concentration of allyl isothiocyanate from all ingredients in the product must

			be no more than 10 mg/kg or 10 mg/L o 0.001%.
3427	MUSTARD SEED OIL	E	Allyl isothiocyanate is a mandatory component of mustard seed oil when the plant part is seed. The concentration of allyl isothiocyanat from all ingredients in the product must be no more than 10 mg/kg or 10 mg/L o 0.001%.
3428	MYOSOTIS ARVENSIS	A, H	
3429	MYRCENE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3430	MYRCENYL ACETATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
3431	MYRICA CERIFERA	A, E, H	
3432	MYRISTIC ACID	Е	
3433	MYRISTIC ALDEHYDE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3434	MYRISTICA FRAGRANS	A, E, H	Safrole is a mandatory component of Myristica fragrans.

Volume	4
--------	---

			<ul> <li>When for internal use then the concentration of safrole in the medicine must be no more than 0.1%.</li> <li>When for topical use then the concentration of safrole in the medicine must be no more than 1%.</li> <li>When the plant preparation is oil or distillate and the concentration in the medicine is more than 50%, the nominal capacity of the container must be no more than 25 millilitres, the medicine must have a restricted flow insert fitted on the container and requires the following warning statement on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or word to that effect).</li> </ul>
3435	MYRISTYL ALCOHOL	E	Only for use in topical medicines for dermal application.
3436	MYRISTYL LACTATE	Е	Only for use in topical medicines for dermal application.
3437	MYRISTYL MYRISTATE	E	Only for use in topical medicines for dermal application.
3438	MYROXYLON BALSAMUM	A, E, H	
3439	MYROXYLON BALSAMUM VAR. PEREIRAE	A, H	
3440	MYRRH	A, H	
3441	MYRRH OIL	A, E, H	
3442	MYRRH RESIN	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>

3443	MYRRHIS ODORATA	A, H	
3444	MYRSINE AFRICANA	A, H	
3445	MYRTENAL	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a flavour.</li><li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li></ul>
3446	MYRTENYL ACETATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3447	MYRTLE OIL	E	Permitted for use only in combination with other permitted ingredients as a fragrance. If used as a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3448	MYRTUS COMMUNIS	A, E, H	
3449	N,N'- BIS(SALICYLIDENE)PROPYLENEDIA MINE	E	N,N'-Bis(salicylidene)propylenediamine must only be included in medicines when in combination with other permitted ingredients as a proprietary excipient formulation in medicines with a dermal route of administration for topical application.
3450	N-(2-(PYRIDIN-2-YL)ETHYL)-P- MENTHANE-3-CARBOXAMIDE	E	<ul> <li>N-(2-(pyridin-2-yl)ethyl)-p-menthane-3- carboxamide must only be included in medicines when in combination with other permitted ingredients as a flavour proprietary excipient formulation.</li> <li>The total concentration of flavour proprietary excipient formulations containing N-(2-(pyridin-2-yl)ethyl)-p- menthane-3-carboxamide must not be more than 5% of the total medicine.</li> </ul>

Volume 4

			The maximum recommended daily dose of the medicine must not provide more than 90 micrograms of N-(2-(pyridin-2- yl)ethyl)-p-menthane-3-carboxamide.
3451	N-BUTYL SULFIDE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3452	N-GLUCONYL ETHANOLAMINE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3453	N-HEXYL 2-BUTENOATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3454	N-NONYL ALCOHOL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3455	NAPHTHALENE	Н	Only for use as an active homoeopathic ingredient.
3456	NARDOSTACHYS CHINENSIS	A, H	

3457	NARINGIN	E	Permitted for use only in combination with other permitted ingredients as a
			flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3458	NASTURTIUM OFFICINALE	A, E, H	
3459	NATURAL FISH OIL	A, E	<ul> <li>When therapeutic indications for this product are made against Vitamin A or colecalciferol (Vitamin D), they are mandatory components of natural fish oil.</li> <li>When for use in topical medicines, the concentration of Vitamin A in the medicine must be no more than 1%.</li> <li>When for internal use, the maximum daily dose must be no more than 3000 micrograms of Retinol Equivalents.</li> <li>When preparations for internal use in adults contain more than 33 microgram of retinol equivalents per dosage unit in divided preparations or per gram of an undivided preparations or per gram of an undivided preparation, the medicine requires the following warning statements on the medicine label:</li> <li>(VITA2) 'WARNING: If you are pregnant - or considering becoming pregnant - do not take vitamin A supplements without consulting your doctor or pharmacist [or words to that effect].' NOTE: Position this warning a the beginning of the directions for use.</li> <li>(VITA4) 'WARNING - When taken in excess of 3000 micrograms retinol equivalents for use.</li> <li>(VITA3) 'The recommended daily amount of vitamin A from all sources is 700 micrograms retinol equivalents for women and 900 micrograms retinol</li> </ul>

## Volume 4

			When for internal use, the maximum recommended daily dose must be no more than 25 micrograms of Vitamin D.
3460	NAUCLEA OFFICINALIS	A, H	
3461	NELUMBO NUCIFERA	A, H	
3462	NELUMBO NUCIFERA FLOWER WAX	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or on damaged skin. The concentration in the medicine must be no more than 0.1%.
3463	NEOHESPERIDIN- DIHYDROCHALCONE	E	<ul> <li>The routes of administration for medicines that contain neohesperidin-dihydrochalcone must be limited to:</li> <li>(a) topical for dermal application; and</li> <li>(b) oral.</li> <li>When used in topical medicines for dermal application:</li> <li>(a) neohesperidin-dihydrochalcone must not be included in medicines intended for use in the eye or on damaged skin; and</li> <li>(b) the concentration of neohesperidin-dihydrochalcone in the medicine must not be more than 0.1%.</li> <li>When used in oral medicines:</li> <li>(a) the concentration in the medicine must not be more than 0.1%; and</li> <li>(b) the following warning statement (or words to that effect) is required on the medicine label:</li> <li>- (NTAKEN3) 'Not to be taken by children under 3 years old'.</li> </ul>
3464	NEOMENTHOL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>

3465	NEOPENTYL GLYCOL DIHEPTANOATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 25%.
			be no more than 25%.
3466	NEOPENTYL GLYCOL DIISOSTEARATE	Е	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 5%.
3467	NEOPENTYL GLYCOL DIOCTANOATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must not be more than 8.1%.
			When the concentration of neopentyl glycol dioctanoate is greater than 5%, the medicine must not be intended for use on damaged skin.
3468	NEOPENTYL GLYCOL DIOCTANOATE/DIDECANOATE	E	Only for use in topical medicines for dermal application.
3469	NEOPICRORHIZA SCROPHULARIIFLORA	A, H	
3470	NEPETA CATARIA	A, H	Pulegone is a mandatory component of Nepeta cataria and must be declared in the application.
			The concentration of pulegone in the medicine must be no more than 4%.
3471	NERAL	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume	4

3472	NERIUM OLEANDER	A, H	The concentration of equivalent dry Nerium oleander in the product must be no more than 1mg/Kg or 1mg/L or 0.0001%.
3473	NEROL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3474	NEROL OXIDE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as part of a flavour or fragrance proprietary excipient formulation.</li> <li>When used in a flavour, the total flavour proprietary excipient formulation in a medicine must be no more than 5%.</li> <li>When used in a fragrance, the total fragrance proprietary excipient formulation in a medicine must be no more than 1%.</li> </ul>
3475	NEROLIDOL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3476	NERONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as part of a fragrance proprietary excipient formulation.</li> <li>The total fragrance proprietary excipient formulation in a medicine must be no more than 1%.</li> </ul>

3477	NERYL ACETATE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3478	NERYL-ISO-BUTYRATE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3479	NICKEL	Н	Only for use as an active homoeopathic ingredient.
3480	NICOTIANA TABACUM	Н	Only for use as an active homoeopathic ingredient.
3481	NICOTINAMIDE	A, E, H	
3482	NICOTINAMIDE ASCORBATE	A, E	
3483	NICOTINAMIDE RIBOSIDE CHLORIDE	А	Ribose is a mandatory component of nicotinamide riboside chloride.
			Only permitted for use in medicines limited to oral routes of administration.
			The maximum recommended daily dose of the medicine must not provide more than 300 mg of nicotinamide riboside chloride.
			The following warning statement (or words to the same effect) is required on the medicine label:
			- (NTAKEN12) 'Not to be taken by children under 12 years old.'
			When the maximum recommended daily dose of the medicine provides greater than 230 mg of nicotinamide riboside

Volume 4	4
----------	---

			<ul> <li>chloride, the following warning statement is required on the medicine label:</li> <li>- (PREG) 'Not recommended for use during pregnancy or lactation'.</li> </ul>
3484	NICOTINIC ACID	A, E	The medicine must contain no more than 100 mg of nicotinic acid per dosage unit
3485	NIGELLA DAMASCENA	A, H	
3486	NIGELLA SATIVA	A, E, H	
3487	NITRIC ACID	E, H	The concentration of nitric acid in the medicine must be no more than 0.5%.
3488	NONADIENOL	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3489	NONANAL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3490	NONANOIC ACID	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.

3491	NONFAT DRY MILK	E, H	
3492	NONIVAMIDE	E	Nonivamide must only be included in medicines when in combination with other permitted ingredients as a flavour or fragrance proprietary excipient formulation.
			The ingredient is not to be included in medicines intended for use in the eye. The total concentration of flavour proprietary excipient formulations containing nonivamide must not be more than 5% of the total medicine.
			The total concentration of fragrance proprietary excipient formulations containing nonivamide must not be more than 1% of the total medicine.
3493	NONOXINOL 10	Е	Only for use in topical medicines for dermal application.
3494	NONOXINOL 12	E	<ul> <li>For use in hand scrub formulations for healthcare professionals only.</li> <li>Only for use in topical medicines for dermal application and not to be include in medicines intended for use in the eye.</li> <li>The concentration in the medicine must be no more than 5%.</li> </ul>
3495	NONOXINOL 5	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li> </ul>
3496	NONOXINOL 9	E	Only for use in topical medicines for dermal application. The concentration in the medicine must be no more than 25%.

Volume 4

3497	NONYL ACETATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
3498	NOOTKATONE	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3499	NOPYL ACETATE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li> </ul>
3500	NORDIHYDROGUAIARETIC ACID	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 0.3%.
3501	NOTOPTERYGIUM FORBESII	A, H	
3502	NOTOPTERYGIUM INCISIUM	A, H	
3503	NUPHAR JAPONICA	A, H	
3504	NUPHAR LUTEA	A, H	
3505	NUTMEG DRY	A, E, H	<ul> <li>Safrole is a mandatory component of Nutmeg Dry.</li> <li>When for internal use then the concentration of safrole from all ingredients in the medicine must be no more than 0.1%.</li> <li>When for topical use then the concentration of safrole from all ingredients in the medicine must be no more than 1%.</li> </ul>

3506	NUTMEG OIL	A, E, H	Safrole is a mandatory component of
			Nutmeg oil.
			When for internal use then the concentration of safrole in the medicine must be no more than 0.1%.
			When for topical use then the concentration of safrole in the medicine must be no more than 1%.
			When the concentration of Nutmeg oil the medicine is more than 50%, the nominal capacity of the container must be no more than 25 mL, the medicine must have a restricted flow insert fitted on the container and requires the following warning statement on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect).
3507	NUTMEG POWDER	A, E, H	Safrole is a mandatory component of Nutmeg powder.
			When for internal use then the concentration of safrole in the medicine must be no more than 0.1%.
			When for topical use then the concentration of safrole in the medicine must be no more than 1%.
3508	NUX VOMICA DRY	A, H	Strychnine (of Strychnos spp.) is a mandatory component of Nux Vomica Dry.
			The concentration of in the medicine must be no more than 1mg/Kg or 1mg/ or 0.0001%.
3509	NUX VOMICA POWDER	Н	Only for use as an active homoeopathic ingredient.
			Strychnine (of Strychnos spp.) is a mandatory component of Nux vomica powder.
			The concentration in the medicine must be no more than 1mg/Kg or 1mg/L or 0.0001%.

## Volume 4

3510	NYCTANTHES ARBOR-TRISTIS	A, H	When the plant part is leaf:
			a) methyl salicylate is a mandatory component of Nyctanthes arbor-tristis;
			b) not to be included in medicines for us in the eye or on damaged skin;
			c) when used internally, the concentration of methyl salicylate in the
			<ul><li>d) when the concentration of methyl salicylate in a liquid preparation is more</li></ul>
			than 5% and the dosage form is other than spray, the medicine requires child resistant packaging;
			<ul> <li>e) when the concentration of methyl salicylate in a liquid preparation is more than 5% and the dosage form is spray, the medicine does not require child resistant packaging if:</li> </ul>
			- the delivery device is engaged into the container in such a way that prevents it from being readily removed;
			- direct suction through the delivery device results in delivery of no more the one dosage unit; and
			- actuation of the spray device is ergonomically difficult for young children to accomplish;
			f) the following warning statement is required on the medicine label:
			- (METSAL) 'Contains methyl salicylate' (or words to that effect); and
			g) when for use in topical medicines for dermal application:
			<ul><li>i) the concentration of methyl salicylate</li><li>in the medicine must not be more than</li><li>25%</li></ul>
			ii) the following warning statements are required on the medicine label:
			- (PREGNT2) 'Do not use if pregnant o likely to become pregnant' (or words to

		medicine
		Volume 4
		<ul> <li>- (CHILD4) 'Do not use [this product/insert name of product] in children 6 years of age or less';</li> <li>- (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect);</li> <li>- (AVOID) 'Avoid prolonged exposure in the sun' (or words to that effect);</li> <li>iii) if the concentration of methyl salicylate in the medicine is greater than 1%, the following warning statement is required on the medicine label:</li> <li>- (IRRIT) 'If irritation develops, discontinue use'.</li> </ul>
NYLON	Е	Only for use in topical medicines for dermal application.
NYLON 6/12	Е	Only for use in topical medicines for dermal application.
NYLON-12	Е	Only for use in topical medicines for dermal application.
NYMPHAEA ALBA	A, E, H	
NYMPHAEA CAERULEA	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine to be no more than 0.3%. Only for use in liquid extracts where the plant part is the flower and the solvent in 100% water.
ΝΥΜΡΗΑΕΑ ΟΠΟΡΑΤΑ	АН	

3511

3512

3513

3514 3515

 3516
 NYMPHAEA ODORATA
 A, H

 3517
 OAK CHIPS EXTRACT
 E
 Permitted for use only in combination with other permitted ingredients as a flavour.

 If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
 If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume	4
--------	---

3518	OAKMOSS ABSOLUTE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3519	OAT	E, H	Only for use as an active homoeopathic or excipient ingredient.
			Gluten is a mandatory component of Oat when the route of administration is other than topical and mucosal.
3520	OAT BRAN	E	Gluten is a mandatory component of Oat bran when the route of administration is other than topical and mucosal.
3521	OATMEAL COLLOIDAL	A, E	Gluten is a mandatory component of Oatmeal colloidal when the route of administration is other than topical and mucosal.
3522	OCIMENE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3523	OCIMENYL ACETATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3524	OCIMUM BASILICUM	A, E, H	When the plant preparation is oil or distillate, Methyl chavicol, eugenol, methyleugenol and cineole are mandatory components of Ocimum basilicum.The concentration of methyleugenol in the medicine must not exceed 1%.When the concentration of Methyl chavicol in the medicine is more than
			5%, the nominal capacity of the container must be no more than 25 millilitres.
			When the concentration of Methyl chavicol in the medicine is more than 5% and the nominal capacity of the containe is 25 millilitres or less, a restricted flow insert must be fitted on the container, an requires the following warning statemen on the medicine label:
			- (CHILD) 'Keep out of reach of children' (or words to that effect). When the concentration of cineole OR eugeno in the preparation is more than 25%, the nominal capacity of the container must not be more than 25 millilitres and the following warning statements must be included on the medicine label:
			<ul> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>- (NTAKEN) 'Not to be taken'.</li> </ul>
			When the concentration of cineole OR eugenol in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres, the medicine must have a child resistant closure and restricted flow insert fitted on the container.
			When the concentration of cineole OR eugenol in the preparation is more than 25% and the nominal capacity of the

Volume 4

			<ul> <li>container is no more than 15 millilitres, the medicine must have a restricted flow insert fitted on the container.</li> <li>When the preparation is for topical use in the mouth, the preparation may not contain more than 5 millilitres of eugenol and the concentration of eugenol in the product must not be greater than 25%.</li> </ul>
3525	OCIMUM KILIMANDSCHARICUM	A, H	Camphor is a mandatory component of Ocimum kilimandscharicum. In solid and semi solid preparations, the concentration of camphor must not be more than 12.5%. In liquid preparations, the nominal capacity of the container must not be more than 25 millilitres. In liquid preparations other than essential oils or distillates, the concentration of camphor must not be more than 2.5%. In essential oil or distillate preparations, if the concentration of camphor is more than 2.5% but less than or equal to 10%, and the nominal capacity of the container is less than 25 millilitres, the medicine must have a restricted flow insert fitted on the container and include the following warning statements on the medicine label: - (CHILD) 'Keep out of reach of children' (or words to that effect); - (NTAKEN) 'Not to be taken'; and - (BABY4) 'Do not apply to infants under 12 months of age except on the advice of a doctor or pharmacist' (or words to that effect). In essential oil or distillate preparations, if the concentration of compton is more
			if the concentration of camphor is more than 10%, and the nominal capacity of the container is less than 15 millilitres, the medicine must have a restricted flow insert fitted on the container and include the following warning statements on the medicine label:

			volume 4
			<ul> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect);</li> <li>- (NTAKEN) 'Not to be taken'; and</li> <li>- (BABY4) 'Do not apply to infants under 12 months of age except on the advice of a doctor or pharmacist' (or words to that effect).</li> <li>In essential oil or distillate preparations, if the concentration of camphor is more than 10%, and the nominal capacity of the container is more than 15 millilitres but less than or equal to 25 millilitres, the medicine must have a restricted flow insert and child resistant closure fitted on the container and include the following warning statements on the medicine label:</li> <li>- (CHILD) 'Keep out of reach of children' (or words to that effect);</li> <li>- (NTAKEN) 'Not to be taken'; and</li> <li>- (BABY4) 'Do not apply to infants</li> </ul>
			under 12 months of age except on the advice of a doctor or pharmacist' (or words to that effect).
3526	OCIMUM MINIMUM	A, H	
3527	OCIMUM TENUIFLORUM	A, H	<ul> <li>When the plant part is oil or distillate, eugenol is a mandatory component of Ocimum tenuiflorum.</li> <li>When the concentration of eugenol in the preparation is more than 25%, the nominal capacity of the container must not be more than 25 millilitres and the following warning statements must be included on the medicine label: <ul> <li>(CHILD) 'Keep out of reach of children' (or words to that effect); and</li> <li>(NTAKEN) 'Not to be taken'.</li> </ul> </li> <li>When the concentration of eugenol in the preparation is more than 25% and the nominal capacity of the container is more than 15 millilitres, the medicine must have</li> </ul>

Vo	lume	4

			<ul> <li>a child resistant closure and restricted flow insert fitted on the container.</li> <li>When the concentration of eugenol in the preparation is more than 25% and the nominal capacity of the container is no more than 15 millilitres, the medicine must have a restricted flow insert fitted on the container.</li> <li>When the preparation is for topical use in the mouth, the preparation may not contain more than 5 mL of eugenol and the concentration of eugenol in the</li> </ul>
3528	OCOTEA ODORIFERA	А, Н	<ul> <li>product must not be greater than 25%.</li> <li>Safrole is a mandatory component of Ocotea odorifera.</li> <li>When for internal use then the concentration of safrole in the medicine must be no more than 0.1%.</li> </ul>
3529	OCTACOSANOL	E	When for topical use then the concentration of safrole in the medicine must be no more than 1%.
3530	OCTADECANAL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3531	OCTADECENE/MA COPOLYMER	Е	Only for use in topical medicines for dermal application.
3532	OCTAHYDRO-4,7-METHANO-3AH- INDENE-3A-CARBOXYLIC ACID, ETHYL ESTER	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>

3533	OCTAHYDROCOUMARIN	Е	Permitted for use only in combination with other permitted ingredients as a
			fragrance. If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.
3534	OCTAN-1-OL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3535	OCTANAL DIMETHYL ACETAL	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3536	OCTANOHYDROXAMIC ACID	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 0.5%.
3537	OCTANOIC ACID	A, E	When for topical use, the concentration in the medicine must be no more than 2% (w/w).
			When for excipient use, permitted for us only in combination with other permitted ingredients as part of a flavour or fragrance proprietary excipient formulation.

Volume 4

			When used in a flavour, the total flavour proprietary excipient formulation in a medicine must be no more than 5%. When used in a fragrance, the total fragrance proprietary excipient formulation in a medicine must be no more than 1%.
3538	OCTENE-1	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as part of a fragrance proprietary excipient formulation.</li> <li>The total fragrance proprietary excipient formulation in a medicine must be no</li> </ul>
3539	OCTOCRYLENE	A	more than 1%.         Only for use as an active ingredient in
3339	OCTOCKTLENE	A	sunscreens for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must not be more than 10%.
			When used in primary sunscreen products, the following warning statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).
3540	OCTOXINOL 10	E	Only for use in topical medicines for dermal application.
3541	OCTYL ACETATE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.

3542	OCTYL CROTONATE	E	Octyl crotonate must only be included in medicines when in combination with other permitted ingredients as a fragrance proprietary excipient formulation.
			The total concentration of the fragrance proprietary excipient formulation containing octyl crotonate must not be more than 1% of the total medicine.
3543	OCTYL HYDROXYSTEARATE	E	Only for use in topical medicines for dermal application.
3544	OCTYL ISOBUTYRATE	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3545	OCTYL ISONONANOATE	E	Only for use in topical medicines for dermal application.
3546	OCTYL METHOXYCINNAMATE	A	Only for use as an active ingredient in sunscreens for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must not be more than 10%.
			When used in primary sunscreen products, the following warning statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).
3547	OCTYL PALMITATE	E	Only for use in topical medicines for dermal application.

Volume 4

3548	OCTYL SALICYLATE	A	Only for use as an active ingredient in sunscreens for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must not be more than 5%.
			When used in primary sunscreen products, the following warning statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure in the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing - hats and eyewear when exposed to the sun' (or words to this effect).
3549	OCTYL STEARATE	E	Only for use in topical medicines for dermal application.
3550	OCTYLBICYCLOHEPTENEDICARBO XIMIDE	Е	Only for use in topical medicines for dermal application.
			The total concentration of octylbicycloheptenedicarboximide in the medicine must not be more than 10%.
3551	OCTYLDODECANOL	E	Only for use in topical medicines for dermal application.
3552	OCTYLDODECETH-25	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 5%.
			Residual levels of 1,4-dioxane and ethylene oxide (and related substances) are to be kept below the level of detection.
3553	OCTYLDODECYL CITRATE CROSSPOLYMER	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye or on damaged skin.
			The concentration in the medicine must be no more than 12%.

3554	OCTYLDODECYL NEOPENTANOATE	E	Only for use in topical medicines for dermal application.
3555	OCTYLDODECYL STEARATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye. The concentration in the medicine must be no more than 2%.
3556	OCTYLDODECYL XYLOSIDE	E	Only for use in topical medicines for dermal application and not to be included in medicines for use in the eye or on damaged skin. The concentration in the medicine must be no more than 1.5%.
3557	OENANTHATE	E	<ul><li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li><li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li></ul>
3558	OENANTHE AQUATICA	Н	Only for use as an active homoeopathic ingredient. The maximum recommended daily dose must be no more than 1mg of the equivalent dry herbal material.
3559	OENANTHE CROCATA	А, Н	The maximum recommended daily dose must be no more than 1mg of the equivalent dry herbal material.
3560	OENOTHERA BIENNIS	A, E, H	
3561	OENOTHERA STRICTA	A, H	
3562	OKOUBAKA AUBREVILLEI	A, H	
3563	OLDENLANDIA DIFFUSA	A, E, H	
3564	OLEA EUROPAEA	A, E, H	
3565	OLEIC ACID	Е	

Volume	4
--------	---

3566	OLETH-10	E	Only for use in topical medicines for dermal application.
3567	OLETH-2	E	Only for use in topical medicines for dermal application.
			Dioxane and Ethylene oxide are mandatory components of Oleth-2.
			The concentration of Dioxane in the medicine must be no more than 10 mg/kg or 10 mg/L or 0.001%.
			The concentration of Ethylene oxide in the medicine must be no more than 1 mg/kg or 1 mg/L or 0.0001%.
3568	OLETH-20	E	Only for use in topical medicines for dermal application.
3569	OLETH-3	E	Only for use in topical medicines for dermal application.
3570	OLETH-3 PHOSPHATE	E	Only for use in topical medicines for dermal application and not to be included in medicines intended for use in the eye.
			The concentration in the medicine must be no more than 0.12%.
3571	OLETH-5	E	Only for use in topical medicines for dermal application.
3572	OLEYL ALCOHOL	E	Only for use in topical medicines for dermal application.
3573	OLIBANUM OIL	A, E, H	
3574	OLIVE OIL	A, E, H	
3575	OMEGA-3 FISH OIL PHYTOSTEROL ESTERS	A	The medicine requires the following warning statement on the medicine label: - (PREGNT) 'Not recommended for use by pregnant and lactating women (or words to that effect).'
3576	OMEGA-3-ACID ETHYL ESTERS 60	A	Docosahexaenoic acid, docosapentaenoic acid and eicosapentaenoic acid are

			Volume 4
			mandatory components of omega-3-acid ethyl esters 60.
			Only permitted for use in medicines that are for oral routes of administration.
			The maximum recommended daily dose of the medicine must not provide more than 3750 milligrams of docosahexaenoic acid, docosapentaenoic acid and eicosapentaenoic acid combined. The following warning statements are required on the medicine label: - (PREGNT) 'Not recommended for use by pregnant and lactating women' (or words to that effect); - (ACOAG) 'Individuals taking anticoagulants should seek medical advice before taking this product' (or words to that effect); - (CHILD3) 'Use in children under 12 years is not recommended'; - (FOOD) 'To be taken with food' (or
			words to that effect).
3577	OMEGA-3-ACID ETHYL ESTERS 90	A	<ul> <li>Only for use in oral medicines.</li> <li>The maximum recommended daily dose of the medicine must not provide more than:</li> <li>a) 4000 mg of omega-3-acid ethyl esters 90; and</li> <li>b) 3750 mg EPA, DHA and DPA combined, when used alone or in combination with other sources of omega-3 fatty acids.</li> </ul>
			The following warning statements (or words to the same effect) are required on the medicine label:
			- (ACOAG) 'Individuals taking anticoagulants should seek medical advice before taking this product.'
			<ul><li> (FOOD) 'To be taken with food.'</li><li> (PREG) 'Not recommended for use during pregnancy or lactation.'</li></ul>

## Volume 4

			- (CHILD3) 'Use in children under 12 years is not recommended.'
3578	ONION	E	
3579	ONION OIL	A, H	
3580	ONONIS SPINOSA	A, E, H	
3581	ONOPORDUM ACANTHIUM	A, H	
3582	ONOSMODIUM VIRGINIANUM	A, H	
3583	OPHIOPOGON JAPONICUS	A, H	
3584	OPOPANAX CHIRONIUM	A, E, H	<ul> <li>When used as an excipient, permitted for use only in combination with other permitted ingredients as part of a flavour or a fragrance proprietary excipient formulation.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3585	OPOPANAX OIL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3586	OPUNTIA FICUS-INDICA	A, H	
3587	ORANGE	E	
3588	ORANGE FLOWER ABSOLUTE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.

3589	ORANGE FLOWER OIL	A, E, H	<ul> <li>When used internally, oxedrine is a mandatory component of orange flower oil.</li> <li>The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.</li> </ul>
3590	ORANGE JUICE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>
3591	ORANGE JUICE OIL	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3592	ORANGE OIL	A, E, H	When used internally, oxedrine is a mandatory component of orange oil. The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
3593	ORANGE OIL BITTER	E	<ul> <li>Orange oil bitter must only be included in medicines when in combination with other permitted ingredients as a flavour or fragrance proprietary excipient formulation.</li> <li>The total concentration of flavour proprietary excipient formulations containing orange oil bitter must not be more than 5% of the total medicine.</li> </ul>
			The total concentration of fragrance proprietary excipient formulations

Volume	4
--------	---

			<ul> <li>containing orange oil bitter must not be more than 1% of the total medicine.</li> <li>The warning statement (SENS)</li> <li>'Application to skin may increase sensitivity to sunlight' or words to that effect must be included on the medicine label unless the medicine is:</li> <li>a) for internal use;</li> <li>b) in preparations containing 1.4% or less of orange oil bitter; or</li> </ul>
			c) for use in soaps or bath or shower ge that are washed off the skin.
3594	ORANGE OIL BITTER COLDPRESSED	A, E, H	When used internally, oxedrine is a mandatory component of orange oil bitter coldpressed.
			The quantity of oxedrine in the maximum recommended daily dose mube no more than 30 milligrams.
			The warning statement (SENS) 'Application to skin may increase sensitivity to sunlight' (or words to that effect) must be included on the medicin label unless the medicine is:
			a) for internal use; or
			b) in preparations containing 1.4% or less of orange oil bitter coldpressed; or
			c) for use in soaps or bath or shower ge that are washed off the skin.
3595	ORANGE OIL COLD PRESSED	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3596	ORANGE OIL DISTILLED	А, Е, Н	When used internally, oxedrine is a mandatory component of orange oil distilled.

			The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
3597	ORANGE OIL SWEET	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.</li> </ul>
3598	ORANGE OIL TERPENELESS	A, E, H	When used internally, oxedrine is a mandatory component of orange oil terpeneless.The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.
3599	ORANGE PEEL	E	Permitted for use only in combination with other permitted ingredients as a flavour. If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3600	ORANGE PEEL DRIED BITTER	A, E, H	<ul> <li>When used internally, oxedrine is a mandatory component of orange peel dried bitter.</li> <li>The quantity of oxedrine in the maximum recommended daily dose must be no more than 30 milligrams.</li> </ul>
3601	ORANGE PEEL OIL SWEET TERPENELESS	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.</li> <li>If used in a flavour the total flavour concentration in a medicine must be no more than 5%.</li> </ul>

Volume 4

			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3602	ORANGE ROUGHY OIL	E	Only for use in topical medicines for dermal application.
3603	ORIGANUM MAJORANA	A, H	<ul> <li>Beta-arbutin is a mandatory component of Origanum majorana.</li> <li>When for oral use, the maximum recommended daily dose must not provide more than 500 mg of beta-arbutin.</li> <li>When for dermal application exclusively to the face: <ul> <li>a) the concentration of beta-arbutin in the medicine must not be more than 7%;</li> <li>b) hydroquinone is a mandatory component; and</li> <li>c) the concentration of hydroquinone must not be more than 10 mg/kg or 10 mg/L or 0.001%.</li> <li>When for use other than oral or dermal application exclusively to the face, the concentration of beta-arbutin in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%.</li> <li>When for use other than oral or dermal application exclusively to the face, the concentration of beta-arbutin in the medicine must not be more than 10 mg/kg or 10 mg/L or 0.001%.</li> <li>When the plant preparation is oil or distillate, and the concentration of Origanum majorana oil or distillate within the medicine is more than 50%:</li> <li>a) the nominal capacity of the container must not be more than 50 mL;</li> <li>b) a restricted flow insert must be fitted on the container; and</li> <li>c) the following warning statement is required on the label: <ul> <li>(CHILD) 'Keep out of reach of children' (or words to that effect).</li> </ul> </li> </ul></li></ul>
3604	ORIGANUM OIL	E	Permitted for use only in combination with other ingredients as a fragrance.

			If used as a fragrance the total concentration in the medicine must be no more than 1%.
3605	ORIGANUM OIL SPANISH	A, E, H	
3606	ORIGANUM VULGARE	A, E, H	
3607	ORNITHINE	A, E	
3608	ORNITHINE ASPARTATE	A, E	
3609	ORNITHINE MONOHYDROCHLORIDE	A, E	
3610	ORNITHOGALUM UMBELLATUM	A, H	
3611	OROSTACHYS FIMBRIATA	A, H	
3612	OROXYLUM INDICUM	A, H	
3613	ORRIS	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3614	ORRIS CONCRETE	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3615	ORRIS ROOT EXTRACT	E	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3616	ORRIS ROOT OIL	A, E, H	
3617	ORRIS ROOT RESIN	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.

Volume 4

## Volume 4

3618	ORTHO-TERT-BUTYLCYCLOHEXYL ACETATE	E	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
			If used in a fragrance the total fragrance concentration in a medicine must be no more 1%.
3619	ORTHOSIPHON ARISTATUS	A, H	
3620	ORYZA SATIVA	A, E, H	
3621	ORYZANOL	Е	
3622	OSBECKIA CHINENSIS	A, H	
3623	OSMANTHUS ABSOLUTE	Е	Permitted for use only in combination with other permitted ingredients as a flavour or a fragrance. If used in a flavour the total flavour
			concentration in a medicine must be no more than 5%.
3624	OSMANTHUS FRAGRANS	Е	Permitted for use only in combination with other permitted ingredients as a flavour.
			If used in a flavour the total flavour concentration in a medicine must be no more than 5%.
3625	OTTELIA ALISMOIDES	A, H	
3626	OXACYCLOHEPTADEC-11-EN-2-ONE	E	<ul> <li>Permitted for use only in combination with other permitted ingredients as a fragrance.</li> <li>If used in a fragrance the total fragrance concentration in a medicine must be no more than 1%.</li> </ul>
3627	OXACYCLOHEXADECAN-2-ONE	E	Only for use in topical medicines for dermal application.

3628	OXACYCLOHEXADECEN-2-ONE	Е	Permitted for use only in combination
			with other permitted ingredients as a
			fragrance.
			If used in a fragrance the total fragrance concentration in a medicine must be no
			more than 1%.
3629	OXALIC ACID	Н	Only for use as an active homoeopathic
			ingredient.
			The total concentration of oxalic acid in the medicine must not be more than 10
			mg/kg or 10 mg/L or 0.001%.
3630	OXALIS ACETOSELLA	A, H	
3631	OXIDISED MAIZE STARCH	Е	Permitted for use only in combination
			with other permitted ingredients as a flavour.
			If used in a flavour the total flavour
			concentration in a medicine must be no
			more than 5%.
3632	OXIDISED TAPIOCA STARCH	E	
3633	OXYBENZONE	А	Only for use as an active ingredient in
			sunscreens for dermal application and not to be included in medicines intended
			for use in the eye.
			The concentration in the medicine must not be more than 10%.
			When used in primary sunscreen
			products, the following warning
			statements are required on the label:
			- (AVOID) 'Avoid prolonged exposure the sun' (or words to this effect); and
			- (SUNPRO) 'Wear protective clothing
			hats and eyewear when exposed to the
			sun' (or words to this effect).
3634	OYSTER	E	The following warning statement is
			required on the medicine label:
			- (MOLLUSC) 'Contains mollusc' or
			'Contains mollusc products'.

Volume 4

3635	OYSTER SHELL	A, E, H	<ul><li>The following warning statement is required on the medicine label:</li><li> (MOLLUSC) 'Contains mollusc' or 'Contains mollusc products'.</li></ul>
------	--------------	---------	--